



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)




PRODUCT INFORMATION



1-Arachidonoyl-d₈-rac-glycerol

Item No. 22694

CAS Registry No.: 2692624-29-4
Formal Name: 5Z,8Z,11Z,14Z-eicosatetraenoic-5,6,8,9,11,12,14,15-d₈ acid, 2,3-dihydroxypropyl ester
Synonym: 1-AG-d₈
MF: C₂₃H₃₀D₈O₄
FW: 386.6
Chemical Purity: ≥95% as a 9:1 mixture of 1-AG and 2-AG
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₈); ≤1% d₀
Supplied as: A solution in acetonitrile
Storage: -80°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

1-Arachidonoyl-d₈-rac-glycerol (1-AG-d₈) is intended for use as an internal standard for the quantification of 1-arachidonoyl glycerol (Item No. 62150) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

1-AG-d₈ is supplied as a solution in acetonitrile. To change the solvent, simply evaporate the acetonitrile under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO purged with an inert gas can be used. The solubility of 1-AG-d₈ in DMSO is approximately 2 mg/ml. 1-AG-d₈ is also miscible in ethanol.

Description

1-AG is a weak cannabinoid (CB₁) receptor agonist and may have other pharmacologic properties.¹ 1-AG is an isomer of 2-AG. 2-AG is chemically unstable and undergoes rapid isomerization to 1-AG (synonymous with 1(3)-AG) both *in vitro* and *in vivo*. 1-AG is a frequent contaminant in synthetic 2-AG preparations, and can markedly reduce their cannabinergic potency.

Reference

1. Sugiura, T., Kodaka, T., Kondo, S., *et al.* Is the cannabinoid CB₁ receptor a 2-arachidonoylglycerol receptor? Structural requirements for triggering a Ca²⁺ transient in NG108-15 cells. *J. Biochem.* **122**(4), 890-895 (1997).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/08/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM