

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



D-Glucosamine-6-sulfate

Item No. 23097

CAS Registry No.:	91674-26-9	
Formal Name:	2-amino-2-deoxy-D-glucose 6-(hydrogen sulfate)	
Synonym:	GlcN-6S	ОН ОН
MF:	C ₆ H ₁₃ NO ₈ S	
FW:	259.2	
Purity:	≥95%	
Supplied as:	A crystalline solid	
Storage:	-20°C	
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

D-Glucosamine-6-sulfate is supplied as a crystalline solid. A stock solution may be made by dissolving the D-glucosamine-6-sulfate in the solvent of choice, which should be purged with an inert gas. D-Glucosamine-6-sulfate is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of D-glucosamine-6-sulfate in these solvents is approximately 2 and 0.5 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of D-glucosamine-6-sulfate can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of D-glucosamine-6-sulfate in PBS, pH 7.2, is approximately 5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

D-Glucosamine-6-sulfate is a naturally occurring glycosaminoglycan.¹ It activates the glmS ribozyme from B. subtilis, a Gram-positive bacterium, when used at a concentration of 200 μ M.² It has been used to form polyvalent dendrimer conjugates that inhibit angiogenesis and endothelial cell proliferation induced by FGF-2 in vitro and prevent scar tissue formation in a rabbit model of glaucoma surgery.³

References

- 1. Foot, M. and Mulholland, M. Classification of chondroitin sulfate A, chondroitin sulfate C, glucosamine hydrochloride and glucosamine 6 sulfate using chemometric techniques. J. Pharm. Biomed. Anal. 38(3), 397-407 (2005).
- 2. Roth, A., Nahvi, A., Lee, M., et al. Characteristics of the glmS ribozyme suggest only structural roles for divalent metal ions. RNA 12(4), 607-619 (2006).
- 3. Shaunak, S., Thomas, S., Gianasi, E., et al. Polyvalent dendrimer glucosamine conjugates prevent scar tissue formation. Nat. Biotechnol. 22(8), 977-984 (2004).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/28/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM