



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION

p,p'-DDE

Item No. 24241

CAS Registry No.: 72-55-9

Formal Name: 1,1'-(2,2-dichloroethenylidene)bis[4-chloro-benzene]

Synonyms: 4,4'-DDE, *p,p'*-Dichlorodiphenyldichloroethylene,
4,4'-Dichlorodiphenyldichloroethylene,
NSC 1153

MF: C₁₄H₈Cl₄


FW: 318.0

Purity: ≥98%

Supplied as: A solid

Storage: -20°C

Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

p,p'-DDE is supplied as a solid. A stock solution may be made by dissolving the *p,p'*-DDE in the solvent of choice, which should be purged with an inert gas. *p,p'*-DDE is slightly soluble in chloroform.

Description

p,p'-DDE is a metabolite and degradation product of the organochlorine pesticide DDT.¹ It accumulates in smallmouth buffalo, channel catfish, and largemouth bass as well as sediment in the Huntsville Spring Branch-Indian Creek tributary system surrounding a DDT manufacturing plant where it is considered a persistent organic pollutant (POP). *p,p'*-DDE inhibits estrogen binding to rainbow trout estrogen receptors (rtERs) with an IC₅₀ value of 8 μM.² It induces concentration-dependent secretion of estradiol by granulosa and theca cell co-cultures isolated from porcine ovarian follicles.³ *p,p'*-DDE (50-100 μM) also decreases ATP levels, the proportion of sperm with high mitochondrial membrane potential, and motility of human sperm *in vitro*.⁴

References

- Garrison, A.W., Cyterski, M., Roberts, K.D., et al. Occurrences and fate of DDT principal isomers/metabolites, DDA, and o,p'-DDD enantiomers in fish, sediment and water at a DDT-impacted superfund site. *Environ. Pollut.* **194**, 224-234 (2014).
- Matthews, J., Celius, T., Halgren, R., et al. Differential estrogen receptor binding of estrogenic substances: A species comparison. *J. Steroid. Biochem. Mol. Biol.* **74(4)**, 223-234 (2000).
- Wójtowicz, A.K., Gregoraszczuk, E.L., Ptak, A., et al. Effect of single and repeated *in vitro* exposure of ovarian follicles to o,p'-DDT and p,p'-DDT and their metabolites. *Pol. J. Pharmacol.* **56(4)**, 465-472 (2004).
- Tavares, R.S., Amaral, S., Paiva, C., et al. *In vitro* exposure to the organochlorine *p,p'*-DDE affects functional human sperm parameters. *Chemosphere* **120**, 443-446 (2015).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.