

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Sphingosine-d₉ (d18:1)

Item No. 24371

Formal Name: (2S,3R,E)-aminooctadec-4-ene-

15,15,16,16,17,17,18,18,18-do-1,3-diol

Synonym: D-erythro-Sphingosine C18-do

MF: $C_{18}H_{28}D_9NO_2$

FW: 308.6


Chemical Purity: ≥98% (Sphingosine)

Deuterium

Incorporation: ≥99% deuterated forms (d₁-d₉); ≤1% d₀

Supplied as: A solid Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Sphingosine-do (d18:1) is intended for use as an internal standard for the quantification of sphingosine (Item No. 10007907) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Sphingosine-do (d18:1) is supplied as a solid. A stock solution may be made by dissolving the sphingosine-d_o (d18:1) in the solvent of choice. Sphingosine-d_o (d18:1) is soluble in organic solvents such as chloroform, ethanol, methanol, and DMSO, which should be purged with an inert gas.

Description

Sphingosine is formed primarily from the breakdown of ceramide. Sphingosine inhibits protein kinase C and phosphatidic acid phosphohydrolase, whereas it activates phospholipase D and diacylglycerol (DAG) kinase. Phosphorylation of sphingosine by sphingosine kinases 1 and 2 (SPHK1, SPHK2) produces sphingosine-1-phosphate, a potent bioactive lipid that exhibits a broad spectrum of biological activities including cell proliferation, survival, migration, cytoskeletal organization, and morphogenesis.²⁻⁴

References

- 1. Hannun, Y.A., Luberto, C., and Argraves, K.M. Enzymes of sphingolipid metabolism: From modular to integrative signaling. Biochemistry 40(16), 4893-4903 (2001).
- 2. Takuwa, Y., Takuwa, N., and Sugimoto, N. The Edg family G protein-coupled receptors for lysophospholipids: Their signaling properties and biological activities. J. Biochem. 131(6), 767-771 (2002).
- Ishii, I., Fukushima, N., Ye, X., et al. Lysophospholipid receptors: Signaling and biology. Annu. Rev. Biochem. 73, 321-354 (2004).
- Kluk, M.J., and Hla, T. Signaling of sphingosine-1-phosphate via the S1P/EDG-family of G-proteincoupled receptors. Biochimica et Biophysica Acta 1582(1-3), 72-80 (2002).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 03/23/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM