

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



COOCH₃

14-methyl Pentadecanoic Acid methyl ester

Item No. 24813

CAS Registry No.: 5129-60-2

Formal Name: 14-methyl-pentadecanoic acid, methyl ester Synonym: Methyl 14-methylpentadecanoate, SFE 17:0

MF: $C_{17}H_{34}O_{2}$ FW: 270.5 **Purity:** ≥98%

Supplied as: A solution in ethanol

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

14-methyl Pentadecanoic acid methyl ester is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as chloroform purged with an inert gas can be used.

Description

14-methyl Pentadecanoic acid methyl ester is a methylated fatty acid methyl ester that has been found in S. zeai sea sponges as the fatty acyl component of zeamide, A. indica leaf extract, and C. vulgaris and H. pluvialis microalgae. 1-3 It is a major component of the vancomycin-induced biofilm produced by vancomycin-resistant S. aureus (VRSA).4 14-methyl Pentadecanoic acid methyl ester has been used as a standard for the quantification of 14-methyl pentadecanoic acid in various foods by GC-MS.⁵

References

- 1. Della Sala, G., Teta, R., Esposito, G., et al. Zeamide, a glycosylinositol phosphorylceramide with the novel core Arap(1β→6)ins motif from the marine sponge Svenzea zeai. Molecules 22(9), 1455 (2017).
- 2. Ravi, S., Shanmugam, B., Subbaiah, G.V., et al. Identification of food preservative, stress relief compounds by GC-MS and HR-LC/Q-TOF/MS; evaluation of antioxidant activity of Acalypha indica leaves methanolic extract (in vitro) and polyphenolic fraction (in vivo). J. Food Sci. Technol. 54(6), 1585-1596 (2017).
- 3. Abdo, S.M., Ali, G.H., and El-Baz, F.K. Potential production of omega fatty acids from microalgae. Int. J. Pharm. Sci. 34(2), 210-215 (2015).
- Mirani, Z.A. and Jamil, N. Role of extra-cellular fatty acids in vancomycin induced biofilm formation by vancomycin resistant Staphylococcus aureus. Pak. J. Pharm. Sci. 26(2), 383-389 (2013).
- 5. Thurnhofer, S. and Vetter, W. Application of ethyl esters and da-methyl esters as internal standards for the gas chromatographic quantification of transesterified fatty acid methyl esters in food. J. Agric. Food Chem. 54(9), 3209-3214 (2006).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/05/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM