



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



7 β -hydroxy Cholesterol-d₇

Item No. 25548

CAS Registry No.: 349553-97-5
Formal Name: (3 β ,7 β)-cholest-5-ene-25,26,26,26,27,27,27-d₇-3,7-diol
MF: C₂₇H₃₉D₇O₂
FW: 409.7
Chemical Purity: \geq 98% (7 β -hydroxy Cholesterol)
Deuterium Incorporation: \geq 99% deuterated forms (d₁-d₇); \leq 1% d₀
Supplied as: A solid
Storage: -20°C
Stability: \geq 4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

7 β -hydroxy Cholesterol-d₇ is intended for use as an internal standard for the quantification of 7 β -hydroxy cholesterol by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated *versus* unlabeled).

7 β -hydroxy Cholesterol-d₇ is supplied as a solid. A stock solution may be made by dissolving the 7 β -hydroxy cholesterol-d₇ in the solvent of choice, which should be purged with an inert gas. 7 β -hydroxy Cholesterol-d₇ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of 7 β -hydroxy cholesterol-d₇ in these solvents is approximately 20, 0.1, and 2 mg/ml, respectively.

Description

7 β -hydroxy Cholesterol is an oxysterol formed by enzymatic and non-enzymatic oxidation of cholesterol.¹ It is the primary oxysterol found in LDL and induces apoptosis and cell death of human umbilical vein endothelial cells (HUVECs) *in vitro* in a concentration-dependent manner. Increased plasma levels of 7 β -hydroxy cholesterol positively correlate with mortality in coronary heart disease in human males.²

References

1. Lizard, G., Monier, S., Cordelet, C., *et al.* Characterization and comparison of the mode of cell death, apoptosis versus necrosis, induced by 7 β -hydroxycholesterol and 7-ketocholesterol in the cells of the vascular wall. *Arterioscler. Thromb. Vasc. Biol.* **19**(5), 1190-1200 (1999).
2. Ziedén, B., Kaminskas, A., Kristenson, M., *et al.* Increased plasma 7 β -hydroxycholesterol concentrations in a population with a high risk for cardiovascular disease. *Artioscler. Thromb. Vasc. Biol.* **19**(4), 967-971 (1999).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/17/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM