

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



N-[1-(2,3-Dioleyloxy)propyl]-N,N,N-trimethylammonium (chloride)

Item No. 25926

CAS Registry No.: 104162-48-3

N,N,N-trimethyl-2,3-bis[(9Z)-9-octadecen-Formal Name:

1-yloxy]-1-propanaminium, monochloride

Synonym:

MF: C₄₂H₈₄NO₂ • CI

FW: 670.6 ≥95% **Purity:**

UV/Vis.: λ_{max} : 225 nm

Supplied as: A solution in ethanol

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



N-[1-(2,3-Dioleyloxy)propyl]-N,N,N-trimethylammonium (DOTMA) (chloride) is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of DOTMA (chloride) in these solvents is approximately 0.5 and 5 mg/ml, respectively.

DOTMA (chloride) is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, the ethanolic solution of DOTMA (chloride) should be diluted with the aqueous buffer of choice. DOTMA (chloride) has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method.

Description

DOTMA is a cationic lipid.¹ It has been used as a component in liposomes that can be used to encapsulate siRNA, microRNAs, and oligonucleotides and for gene transfection in vitro. 1,2

References

- 1. Wang, X., Yu, B., Ren, W., et al. Enhanced hepatic delivery of siRNA and microRNA using oleic acid based lipid nanoparticle formulations. J. Control Release 172(3), 690-698 (2013).
- 2. Huang, Q.-D., Zhong, G.-X., Zhang, Y., et al. Cyclen-based cationic lipids for highly efficient gene delivery towards tumor cells. PLoS One 6(8), e23134 (2011).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 10/19/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM