

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Trimethoprim-d₃

Item No. 26119

CAS Registry No.: 1189923-38-3

Formal Name: 5-(3,5-dimethoxy-4-(methoxy-d₃)benzyl)

pyrimidine-2,4-diamine

MF: $C_{14}H_{15}D_3N_4O_3$

FW: 293.3


Chemical Purity: ≥98% (Trimethoprim)

Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₃); \leq 1% d₀

Supplied as: A solid Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Trimethoprim-d₃ is intended for use as an internal standard for the quantification of trimethoprim (Item No. 16473) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Trimethoprim-d₂ is supplied as a solid. A stock solution may be made by dissolving the trimethoprim-d₂ in the solvent of choice, which should be purged with an inert gas. Trimethoprim-d₃ is slightly soluble in DMSO and methanol.

Description

Trimethoprim is a synthetic antibiotic that inhibits dihydrofolate reductase (DHFR), which is necessary for the synthesis of purines, amino acids, and thymidylic acid. 1,2 Trimethoprim shows prominent selectivity for bacterial DHFR over mammalian DHFR (IC_{50} s = 5 and 30,000 nM, respectively).² Formulations containing trimethoprim, commonly used in combination with sulfamethoxazole to minimize acquired resistance, have been used in the treatment of urinary tract infections.

References

- 1. Lam, T., Hilgers, M.T., Cunningham, M.L., et al. Structure-based design of new dihydrofolate reductase antibacterial agents: 7-(Benzimidazol-1-yl)-2,4-diaminoquinazolines. J. Med. Chem. 57(3), 651-668 (2014).
- 2. Schweitzer, B.I., Dicker, A.P., and Bertino, J.R. Dihydrofolate reductase as a therapeutic target. FASEB J. 4(8), 2441-2452 (1990).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Suyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 09/21/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM