

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Azido-Thalidomide

Item No. 26146

CAS Registry No.: 2098488-36-7

Formal Name: N-(4-azidobutyl)-2-[[2-(2,6-dioxo-3-


> piperidinyl)-2,3-dihydro-1,3-dioxo-1H-isoindol-4-yl]oxy]-acetamide

MF: $C_{19}H_{20}N_6O_6$ FW: 428.4 ≥95% **Purity:**

UV/Vis.: λ_{max} : 219 nm Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Azido-thalidomide is supplied as a crystalline solid. A stock solution may be made by dissolving the azido-thalidomide in the solvent of choice. Azido-thalidomide is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of azido-thalidomide in these solvents is approximately 12 mg/ml.

Azido-thalidomide is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, azido-thalidomide should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Azido-thalidomide has a solubility of approximately 0.11 mg/ml in a 1:8 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Azido-thalidomide is a building block in the synthesis of proteolysis-targeting chimera (PROTAC) technologies. It contains thalidomide, which is a ligand for the E3 ubiquitin ligase cereblon, and can be conjugated to alkynylated ligands for target proteins via click chemistry. Azido-thalidomide has been used in the synthesis of a SirReal-based PROTAC for the degradation of sirtuin 2 (Sirt2) in cells.

Reference

1. Schiedel, M., Herp, D., Hammelmann, S., et al. Chemically induced degradation of Sirtuin 2 (Sirt2) by a proteolysis targeting chimera (PROTAC) based on sirtuin rearranging ligands (SirReals). J. Med. Chem. 61(2), 482-491 (2018).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 01/18/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM