

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



WAY-316606 Item No. 26324

CAS Registry No.: 915759-45-4

Formal Name: 5-(phenylsulfonyl)-N-4-

piperidinyl-2-(trifluoromethyl)-

benzenesulfonamide

MF: $C_{18}H_{19}F_3N_2O_4S_2$

448.5 FW: **Purity:** ≥98% UV/Vis.: λ_{max} : 239 nm Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

WAY-316606 is supplied as a crystalline solid. A stock solution may be made by dissolving the WAY-316606 in the solvent of choice. WAY-316606 is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of WAY-316606 in these solvents is approximately 3 mg/ml.

WAY-316606 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, WAY-316606 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. WAY-316606 has a solubility of approximately 0.1 mg/ml in a 1:10 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

WAY-316606 is an inhibitor of secreted frizzled-related protein 1 (sFRP-1; $IC_{50} = 0.5 \mu M$).¹ It increases Wnt signaling in U2OS osteosarcoma cells (EC₅₀ = 0.65 μ M). WAY-316606 increases total bone area in a murine calvarial organ culture assay when used at concentrations greater than 0.1 nM.²

References

- 1. Moore, W.J., Kern, J.C., Bhat, R., et al. Modulation of Wnt signaling through inhibition of secreted frizzledrelated protein I (sFRP-1) with N-substituted piperidinyl diphenylsulfonyl sulfonamides: Part II. Bioorg. Med. Chem. 18(1), 190-201 (2010).
- 2. Bodine, P.V., Stauffer, B., Ponce-de-Leon, H., et al. A small molecule inhibitor of the Wnt antagonist secreted frizzled-related protein-1 stimulates bone formation. Bone 44(6), 1063-1068 (2009).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/08/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM