

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Cobicistat-d₈ Item No. 26442

Formal Name: thiazol-5-ylmethyl ((2R,5R)-5-((S)-2-

> (3-((2-isopropylthiazol-4-yl)methyl)-3-methylureido)-4-(morpholino-d_o) butanamido)-1,6-diphenylhexan-2-yl)

carbamate

MF: $C_{40}H_{45}D_8N_7O_5S_2$

FW: 784.1


Chemical Purity: ≥98% (Cobicistat)

Deuterium

≥99% deuterated forms (d₁-d₈); ≤1% d₀ Incorporation:

Supplied as: A solid -20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Cobicistat-d₈ is intended for use as an internal standard for the quantification of cobicistat (Item No. 23433) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Cobicistat-d₈ is supplied as a solid. A stock solution may be made by dissolving the cobicistat-d₈ in the solvent of choice. Cobicistat-d₈ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of cobicistat-d_s is approximately 10 mg/ml in ethanol and approximately 20 mg/ml in DMSO and DMF.

Description

Cobicistat is an inhibitor of the cytochrome P450 (CYP) isomer CYP3A (IC₅₀s = 30-285 nM for CYP3A metabolism of various HIV protease inhibitors).1 It is selective for CYP3A over other CYP isomers (IC₅₀ = >25 μ M for CYP1A2, 2C8, 2C9, and 2C19). Cobicistat does not inhibit HIV-1 protease $(IC_{50}^{30} = 30 \mu M)$ or affect HIV replication in MT-2 cells (EC₅₀ = 30 μM). Formulations containing cobicistat have been used to slow the metabolism of concomitantly administered protease inhibitors in the treatment of HIV.2

References

- 1. Xu, L., Liu, H., Murray, B.P., et al. Cobicistat (GS-9350): A potent and selective inhibitor of human CYP3A as a novel pharmacoenhancer. ACS Med. Chem. Lett. 1(5), 209-213 (2010).
- 2. Harris, M., Ganase, B., Watson, B., et al. HIV treatment simplification to elvitegravir/cobicistat/ emtricitabine/tenofovir disproxil fumarate (E/C/F/TDF) plus darunavir: A pharmacokinetic study. AIDS Res. Ther. 14(1), 59 (2017).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/15/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM