

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Octanoyl-L-carnitine (chloride)

Item No. 26558

CAS Registry No.: 54377-02-5

Formal Name: (2R)-3-carboxy-N,N,N-

> trimethyl-2-[(1-oxooctyl)oxy]-1propanaminium, monochloride

Synonyms: C8 Carnitine, L-Carnitine octanoyl,

L-Octanoylcarnitine ester


 $C_{15}H_{30}NO_4 \bullet CI$ MF:

323.9 FW: **Purity:** ≥95%

Supplied as: A crystalline solid

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Octanoyl-L-carnitine (chloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the octanoyl-L-carnitine (chloride) in the solvent of choice. Octanoyl-L-carnitine (chloride) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of octanoyl-L-carnitine (chloride) in ethanol and DMF is approximately 20 mg/ml and approximately 10 mg/ml in DMSO.

Description

Octanoyl-L-carnitine is a medium-chain acylcarnitine and the physiologically active form of octanoyl-DL-carnitine (Item No. 15048). Plasma levels of octanoyl-L-carnitine are elevated in patients with end-stage renal disease on continuous ambulatory peritoneal dialysis (PD) compared with both patients on automated PD and healthy individuals.² Octanoyl-L-carnitine does not undergo hydrolysis in the blood or during sample preparation when used as a standard for the quantification of octanoylcarnitine.¹

References

- 1. Chace, D.H., DiPerna, J.C., Adam, B.W., et al. Errors caused by the use of D,L-octanoylcarnitine for blood-spot calibrators. Clin. Chem. 47(4), 758-760 (2001).
- 2. Di Liberato, L., Arduini, A., Rossi, C., et al. L-Carnitine status in end-stage renal disease patients on automated peritoneal dialysis. J. Nephrol. 27(6), 699-706 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 04/04/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM