

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Sterculic Acid

Item No. 26735

CAS Registry No.: 738-87-4

Formal Name: 2-octyl-1-cyclopropene-1-

octanoic acid

MF: $C_{19}H_{34}O_{2}$ FW: 294.5 **Purity:** ≥98% Supplied as: A liquid Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Sterculic acid is supplied as a liquid. Sterculic acid is soluble in organic solvents such as chloroform, hexane, ethyl ether, and methanol, which should be purged with an inert gas.

Description

Sterculic acid is a cyclopropene fatty acid that has been found in S. foetida. 1 It is an inhibitor of Δ^9 -desaturase that decreases 9(Z)-myristoleic acid (Item No. 9002461) and increases oleic acid (Item No. 90260) and 9(Z),11(E)-conjugated linoleic acid (Item No. 90140) levels in the milk of lactating ewes when administered at a dose of 0.5 g per day.² It also inhibits endoplasmic reticulum stress induced by 7-keto cholesterol (Item No. 16339) in ARPE-19 cells when used at a concentration of 1 µM and prevents the formation of choroidal neovascularization when applied to eyes at concentrations of 0.1 to 10 mM in a rat model of macular degeneration induced by laser injury. Sterculic acid binds to a variety of kinases, including calcium/calmodulin-dependent protein kinase kinase 2 (CAMKK2), mammalian sterile20-related kinase 3 (MST3), and p90 ribosomal S5 kinase 4 (RSK4).3

References

- 1. Huang, J.-D., Amaral, J., Lee, J.W., et al. Sterculic acid antagonizes 7-ketocholesterol-mediated inflammation and inhibits choroidal neovascularization. Biochim. Biophys. Acta. 1821(4), 637-646 (2012).
- 2. Bichi, E., Toral, P.G., Hervás, G., et al. Inhibition of Δ^9 -desaturase activity with sterculic acid: Effect on the endogenous synthesis of cis-9 18:1 and cis-9, trans-11 18:2 in dairy sheep. J. Dairy Sci. 95(9), 5242-5252 (2012).
- 3. Huang, J.-D., Amaral, J., Lee, J.W., et al. 7-Ketocholesterol-induced inflammation signals mostly through the TLR4 receptor both in vitro and in vivo. PLoS One 9(7), e100985 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/29/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM