

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



(R)-α-Lipoic Acid

Item No. 27299

CAS Registry No.: 1200-22-2

Formal Name: (3R)-1,2-dithiolane-3-pentanoic acid

Synonym: (R)-(+)-Lipoic Acid MF:

 $C_8H_{14}O_2S_2$ FW: 206.3 **Purity:** ≥98%

 λ_{max} : 331 nm A crystalline solid UV/Vis.: Supplied as:

-20°C Storage: Stability: ≥2 vears

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

(R)-α-Lipoic acid is supplied as a crystalline solid. A stock solution may be made by dissolving the (R)- α -lipoic acid in the solvent of choice. (R)- α -Lipoic acid is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of $(R)-\alpha$ -lipoic acid in these solvents is approximately 30 mg/ml.

(R)-α-Lipoic acid is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, (R)- α -lipoic acid should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. (R)-α-Lipoic acid has a solubility of approximately 0.25 mg/ml in a 1:8 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

(R)-α-Lipoic acid is the naturally occurring enantiomer of lipoic acid, a cyclic disulfide antioxidant. It acts as a cofactor in 2-oxo dehydrogenase complex reactions, where it is reduced to dihydrolipoic acid (DHLA; Item No. 16372). (R)-α-Lipoic acid (100 mg/kg per day) decreases serum levels of malondialdehyde (MDA) and increases the activity of superoxide dismutase (SOD) and total antioxidative capabilities (T-AOC) in a mouse model of accelerated senescence induced by D-galactose (Item No. 20890).2 It also reduces the escape latency in the Morris water maze compared to both D-galactose and vehicle control groups and rescues decreases in neural progenitor cell proliferation in the hippocampal dentate gyrus.

References

- 1. Biewenga, G.P., Haenen, G.R.M.M., and Bast, A. The pharmacology of the antioxidant lipoic acid. Gen. Pharmacol. 29(3), 315-331 (1997).
- 2. Cui, X., Zuo, P., Zhang, Q., et al. Chronic systemic D-galactose exposure induces memory loss, neurodegeneration, and oxidative damage in mice: Protective effects of R- α -lipoic acid. J. Neurosci. Res. 84(3), 647-654 (2006).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the mater can be found on our website.

Copyright Cayman Chemical Company, 06/24/2019

COOH

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM