

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



BLX3887

Item No. 27391

CAS Registry No.: 934758-70-0

Formal Name: 4,5-dichloro-N-(2-chloro-4-

fluorophenyl)-1H-pyrazole-3-

carboxamide

Synonym: 15-LO-1 Inhibitor 3887


MF: $C_{10}H_5CI_3FN_3O$

FW: 308.5 **Purity:** ≥95%

 λ_{max} : 260 nm A crystalline solid UV/Vis.: Supplied as:

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

BLX3887 is supplied as a crystalline solid. A stock solution may be made by dissolving the BLX3887 in the solvent of choice, which should be purged with an inert gas. BLX3887 is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of BLX3887 in these solvents is approximately 15 and 30 mg/ml, respectively.

BLX3887 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, BLX3887 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. BLX3887 has a solubility of approximately 0.16 mg/ml in a 1:5 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

BLX3887 is an inhibitor of 15-lipoxygenase type 1 (15-LO-1; IC_{50} = 32 nM in a cell-free enzyme assay).¹ It is selective for 15-LO-1 over 15-LO-2, which it does not inhibit, 5-LO (IC_{50} = 472 nM), and 12-LO (IC₅₀ = 3,310 nM). BLX3887 inhibits the production of 15-LO metabolites selectively in eosinophils over neutrophils when used at a concentration of 10 µM. It also inhibits endocytosis in, and the migration of, isolated human peripheral blood mononuclear cell-derived dendritic cells in vitro.

Reference

1. Archambault, A.-S., Turcotte, C., Martin, C., et al. Comparison of eight 15-lipoxygenase (LO) inhibitors on the biosynthesis of 15-LO metabolites by human neutrophils and eosinophils. PLoS One 13(8), e0202424 (2018).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 01/22/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM