



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## Mulberroside A

Item No. 27545

**CAS Registry No.:** 102841-42-9  
**Formal Name:** 3-[(1E)-2-[4-(β-D-glucopyranosyloxy)-2-hydroxyphenyl]ethenyl]-5-hydroxyphenyl, β-D-glucopyranoside  
**MF:** C<sub>26</sub>H<sub>32</sub>O<sub>14</sub>  
**FW:** 568.5  
**Purity:** ≥98%  
**UV/Vis.:** λ<sub>max</sub>: 216, 301, 324 nm  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:** ≥2 years  
**Item Origin:** Plant/Cortex mori



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Mulberroside A is supplied as a crystalline solid. A stock solution may be made by dissolving the mulberroside A in the solvent of choice, which should be purged with an inert gas. Mulberroside A is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of mulberroside A in ethanol is approximately 2 mg/ml and approximately 25 mg/ml in DMSO and DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of mulberroside A can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of mulberroside A in PBS, pH 7.2, is approximately 0.5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

Mulberroside A is a phenol and stilbene glucoside form of oxyresveratrol (Item No. 12028) originally isolated from *M. alba* (mulberry) roots that has diverse biological activities.<sup>1-4</sup> It inhibits pregnane X receptor-mediated P-glycoprotein (P-gp) luciferase reporter activity induced by rifampicin (Item No. 14423) in LS174T cells when used at concentrations ranging from 5 to 20 μM.<sup>1</sup> Mulberroside A decreases expression of TNF-α, IL-1β, and IL-6, inhibits caspase-1 activation, and increases cell viability in an isolated rat cortical neuron model of oxygen-glucose deprivation-induced ischemia and reperfusion injury.<sup>2</sup> Topical administration of mulberroside A (2 and 5% v/v) reduces UVB-induced hyperpigmentation, levels of the melanogenesis enzymes tyrosinase, tyrosinase-related protein 1 (TRP-1), and microphthalmia-associated transcription factor (MITF), and tyrosinase activity in brown guinea pig skin.<sup>3</sup> Mulberroside A (10, 20, and 40 mg/kg) decreases serum levels of uric acid, creatinine, and urea nitrogen, reduces renal vacuolar and granular degeneration, and increases fractional excretion of uric acid and urinary urate excretion in hyperuricemic mice.<sup>4</sup>

### References

1. Li, Y., Huang, L., Sun, J., et al. *J. Biochem. Mol. Toxicol.* (2016).
2. Wang, C.-P., Zhang, L.-Z., Li, G.-C., et al. *J. Neurosci. Res.* **92**(7), 944-954 (2014).
3. Park, K.-T., Kim, J.-K., Hwang, D., et al. *Food Chem. Toxicol.* **49**(12), 3038-3045 (2011).
4. Wang, C.-P., Wang, Y., Wang, X., et al. *Planta Med.* **77**(8), 786-794 (2011).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/03/2019

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM