



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

Resorufin β -D-Galactopyranoside

Item No. 28015

CAS Registry No.: 95079-19-9
 Formal Name: 7-(β -D-galactopyranosyloxy)-3H-phenoxazin-3-one
 MF: $C_{18}H_{17}NO_8$
 FW: 375.3
 Purity: $\geq 98\%$
 UV/Vis.: λ_{max} : 245, 458 nm
 Ex./Em. Max: 570/580 nm
 Supplied as: A solid
 Storage: -20°C
 Stability: ≥ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Resorufin β -D-galactopyranoside is supplied as a solid. A stock solution may be made by dissolving the resorufin β -D-galactopyranoside in the solvent of choice, which should be purged with an inert gas. Resorufin β -D-galactopyranoside is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of resorufin β -D-galactopyranoside in these solvents is approximately 1, 25, and 2 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of resorufin β -D-galactopyranoside can be prepared by directly dissolving the solid in aqueous buffers. The solubility of resorufin β -D-galactopyranoside in PBS, pH 7.2, is approximately 0.3 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Resorufin β -D-galactopyranoside is a fluorogenic substrate for β -galactosidase.¹ Upon enzymatic cleavage by β -galactosidase, the fluorescent moiety resorufin is released and its fluorescence can be used to quantify β -galactosidase activity. Resorufin displays excitation/emission maxima of 570/580 nm, respectively.

Reference

- Hofmann, J. and Sernetz, M. Immobilized enzyme kinetics analyzed by flow-through microfluorimetry: Resorufin- β -D-galactopyranoside as a new fluorogenic substrate for β -galactosidase. *Analytica Chimica Acta* **163**, 67-72 (1984).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/22/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
 ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
 [734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
 WWW.CAYMANCHEM.COM