

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Nevirapine-d₄

Item No. 28093

CAS Registry No.: 1051418-95-1

Formal Name: 11-cyclopropyl-2,2,3,3-d₄-5,11-dihydro-

4-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]

diazepin-6-one

Synonyms: NVP-d₄

MF: $C_{15}H_{10}D_4N_4O$

FW: 270.3


Chemical Purity: ≥98% (Nevirapine)

Deuterium

Incorporation: ≥99% deuterated forms (d_1-d_4) ; ≤1% d_0

Supplied as: A solid -20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Nevirapine-d₄ is intended for use as an internal standard for the quantification of nevirapine (Item No. 15117) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Nevirapine- d_{Δ} is supplied as a solid. A stock solution may be made by dissolving the nevirapine- d_{Δ} in the solvent of choice. Nevirapine- d_4 is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of nevirapine-d₁ in these solvents is approximately 2.5 and 5 mg/ml, respectively.

Description

Nevirapine is a non-nucleoside reverse transcriptase inhibitor (NNRTI).¹ It binds to HIV-1 reverse transcriptase and inhibits RNA plus-strand initiation (IC $_{50}$ = 0.45 μ M). Nevirapine prevents seroconversion and viremia in a chimpanzee model of HIV-1 infection.² Formulations containing nevirapine have been used in combination therapy for the treatment of HIV-1 infection.

References

- 1. Grobler, J.A., Dornadula, G., Rice, M.R., et al. HIV-1 reverse transcriptase plus-strand initiation exhibits preferential sensitivity to non-nucleoside reverse transcriptase inhibitors in vitro. J. Biol. Chem. 282(11), 8005-8010 (2007).
- Grob, P.M., Cao, Y., Muchmore, E., et al. Prophylaxis against HIV-1 infection in chimpanzees by nevirapine, a nonnucleoside inhibitor of reverse transcriptase. Nat. Med. 3(6), 665-670 (1997).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 07/22/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM