

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



cis-6-Hexadecenoic Acid-d₁₀

Item No. 28227

Formal Name: (Z)-hexadec-6-enoic-8,8,9,9,10,10,11,11,12,

12,13,13,14,14,15,15,16,16,16-d₁₉ acid

Synonyms: C16:1(6Z)-d₁₉, C16:1Δ6-d₁₉,

C16:1ω10c-d₁₉, Sapienic Acid-d₁₉

 $C_{16}H_{11}D_{19}O_2$ MF:

FW: 273.5

Chemical Purity: ≥98% (cis-6-Hexadecenoic Acid)


Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₁₉); \leq 1% d₀

Supplied as: A solution in ethanol

Storage: -20°C Stability: ≥1 year

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

cis-6-Hexadecenoic acid-d₁₉ is intended for use as an internal standard for the quantification of cis-6-hexadecenoic acid (Item No. 9001845) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Description

cis-6-Hexadecenoic acid is a monounsaturated fatty acid and is one of the primary fatty acids in human skin. 2 cis-6-Hexadecenoic acid levels are increased in isolated sebum from the face and back of patients with acne.² In contrast, levels are decreased in the non-lesional skin and isolated sebum of atopic dermatitis patients, which correlates with an increase in S. aureus in the sebum. It is active against S. aureus in vitro when used at a concentration of 5 μ g/ml at pH 5.5.3 cis-6-Hexadecenoic acid disrupts membrane integrity, the proton motive force, increases membrane fluidity, and inhibits the electron transport chain in S. aureus.

References

- 1. Takigawa, H., Nakagawa, H., Kuzukawa, M., et al. Deficient production of hexadecenoic acid in the skin is associated in part with the vulnerability of atopic dermatitis patients to colonization by Staphylococcus aureus. Dermatology **211(3)**, 240-248 (2005).
- 2. Li, W.-H., Zhang, Q., Flach, C.R., et al. In vitro modeling of unsaturated free fatty acid-mediated tissue impairments seen in acne lesions. Arch. Dermatol. Res. 309(7), 529-540 (2017).
- 3. Cartron, M.L., England, S.R., Chiriac, A.I., et al. Bactericidal activity of the human skin fatty acid cis-6-hexadecanoic acid on Staphylococcus aureus. Antimicrob. Agents Chemother. 58(7), 3599-3609 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 08/16/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM