

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



.COOH

Arachidic Acid-d₂

Item No. 28083

CAS Registry No.: 232600-70-3

Formal Name: eicosanoic-2,2-d2 acid

Synonyms: Arachic Acid-d₂, Arachidate-d₂,

Eicosanoate-d₂, Icosanoic Acid-d₂

MF: $C_{20}H_{38}D_2O_2$ FW: 314.5

Chemical Purity: ≥95% (Arachidic Acid)

Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₂); \leq 1% d₀

Supplied as: A crystalline solid

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Arachidic acid-d₂ is intended for use as an internal standard for the quantification of arachidic acid (Item Nos. 9000339 | 21906) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Arachidic acid-d₂ is supplied as a crystalline solid. A stock solution may be made by dissolving the arachidic acid-d₂ in the solvent of choice. Arachidic acid-d₂ is soluble in organic solvents such as ethanol and dimethyl formamide, which should be purged with an inert gas. The solubility of arachidic acid-d2 in these solvents is approximately 0.1 and 2 mg/ml, respectively.

Description

Arachidic acid is a long-chain saturated fatty acid that has been found in peanut butter and anaerobic fungi. ^{1,2} It inhibits rabbit neutrophil aggregation induced by N-formyl-methionyl-leucyl-phenylalanine (fMLP; Item No. 21495) when used at a concentration of 5 μ M.³ Formulations containing arachidic acid have been used as surfactants in the manufacture of soaps and cosmetics.

References

- 1. Negoita, M., Mihai, A.L., Adascalului, A., et al. Comparison of the fatty acid composition of peanut butter by applying different fat extraction procedures. Rev. Chim. (Bucharest) 69(11), 3023-3032 (2018).
- Koppová, I., Novotná, Z., Štrosová, L., et al. Analysis of fatty acid composition of anaerobic rumen fungi. Folia Microbiol. (Praha) 53(3), 217-220 (2008).
- Naccache, P.H., Moiski, T.F., Volpi, M., et al. Modulation of rabbit neutrophil aggregation and degranulation by free fatty acids. J. Leukoc. Biol. 36(3), 333-340 (1984).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 01/08/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM