



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



IT1t (hydrochloride)

Item No. 28434

CAS Registry No.: 1092776-63-0

Formal Name: N,N'-dicyclohexyl-carbamimidothioic acid,
(5,6-dihydro-6,6-dimethylimidazo[2,1-b]
thiazol-3-yl)methyl ester, dihydrochloride

MF: C₂₁H₃₄N₄S₂ • 2HCl

FW: 479.6


Purity: ≥95%

UV/Vis.: λ_{max}: 233 nm

Supplied as: A crystalline solid

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

IT1t (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the IT1t (hydrochloride) in the solvent of choice, which should be purged with an inert gas. IT1t (hydrochloride) is soluble in the organic solvent DMSO.

Description

IT1t is a chemokine (C-X-C motif) receptor 4 (CXCR4) antagonist (IC₅₀s = 8 and 11 nM for the human and rat receptors, respectively).¹ It is selective for CXCR4 over human ether-a-go-go-related gene potassium channels (hERG/K_v11.1; IC₅₀ = 13,240 nM). IT1t inhibits calcium mobilization induced by chemokine (C-X-C-motif) ligand 12 (CXCL12) in CEM cells (IC₅₀ = 1.1 nM) and decreases CXCL12-induced migration of Jurkat cells (IC₅₀ = 79.1 nM).^{1,2} It inhibits replication of the HIV-1 strain NL4-3 in MT-4 cells and isolated human peripheral blood mononuclear cells (PBMCs) stimulated with phytohemagglutinin (PHA; IC₅₀s = 14.2 and 19 nM, respectively).² IT1t (20 μM) reduces tumor growth in an MDA-MB-231-B zebrafish xenograft model.³

References

- Thoma, G., Streiff, M.B., Kovarik, J., et al. Orally bioavailable isothioureas block function of the chemokine receptor CXCR4 in vitro and in vivo. *J. Med. Chem.* **51**(24), 7915-7920 (2008).
- Van Hout, A., D'huys, T., Oeyen, M., et al. Comparison of cell-based assays for the identification and evaluation of competitive CXCR4 inhibitors. *PLoS One* **12**(4), e0176057 (2017).
- Tulotta, C., Stefanescu, C., Beletkaia, E., et al. Inhibition of signaling between human CXCR4 and zebrafish ligands by the small molecule IT1t impairs the formation of triple-negative breast cancer early metastases in a zebrafish xenograft model. *Dis. Model Mech.* **9**(2), 141-153 (2016).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/29/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM