

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



trans-Zeatin-d₅

Item No. 28534

Formal Name: 2E-methyl-d₃-4-(9H-purin-6-ylamino)-

but-2-en-1,1-d₂-1-ol

MF: $C_{10}H_8D_5N_5O$

FW:


Chemical Purity: ≥98% (trans-Zeatin)

Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₅); \leq 1% d₀

Supplied as: A solid Storage: -20°C Stability: ≥2 years Item Origin: Synthetic

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

trans-Zeatin-d₅ is intended for use as an internal standard for the quantification of trans-zeatin (Item No. 13226) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

trans-Zeatin-d_s is supplied as a solid. A stock solution may be made by dissolving the trans-zeatin-d_s in the solvent of choice, which should be purged with an inert gas. trans-Zeatin-d₅ is soluble in methanol and DMSO.

Description

trans-Zeatin is a cytokinin plant growth regulator with antioxidant and neuroprotective activities.¹⁻⁴ It binds to the cytokinin receptor Arabidopsis histidine kinase 3 (AHK3) with a K_D value of 1.3 nM.3 trans-Zeatin increases chlorophyll levels in etiolated Cucumus sativus cotyledons in a concentration-dependent manner.¹ It increases callus growth and shoot formation in N. tabacum calluses when used at concentrations of 5 and 50 μM.² trans-Zeatin (25-100 μM) reduces production of reactive oxygen species (ROS) induced by amyloid β (25-35) (Aβ25-35) in PC12 cells.⁴ It reduces scopolamine-induced spontaneous alternations in the Y-maze, indicating enhanced spatial memory, in mice when administered at doses of 1.5, 3, and 4.5 mg/kg per day.

References

- 1. Fletcher, R.A. and McCullagh, D. Cytokinin-induced chlorophyll formation in cucumber cotyledons. Planta **101(1)**, 88-90 (1971).
- Yamada, Y., Sekiya, J., and Koshimizu, K. Cytokinin-induced shoot formation. Phytochemistry 11(3), 1019-1021 (1972).
- 3. Romanov, G.A., Lomin, S.N., and Schmülling, T. Biochemical characteristics and ligand-binding properties of Arabidopsis cytokinin receptor AHK3 compared to CRE1/AHK4 as revealed by a direct binding assay. J. Exp. Bot. 57(15), 4051-4058 (2006).
- 4. Choi, S.J., Jeong, C.-H., Choi, S.-G., et al. Zeatin prevents amyloid β-induced neurotoxicity and scopolamine-induced cognitive deficits. J. Med. Food 12(2), 271-277 (2009).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 01/07/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM