

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



CU-115

Item No. 29815

CAS Registry No.: 2471982-20-2

Formal Name: N-[4-[3,5-bis(trifluoromethyl)

phenoxylphenyll-2-fluoro-6-iodo-

benzamide

MF: $C_{21}H_{11}F_7INO_2$

569.2 FW: **Purity:** ≥98%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 vears

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

CU-115 is supplied as a crystalline solid. A stock solution may be made by dissolving the CU-115 in the solvent of choice, which should be purged with an inert gas. CU-115 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of CU-115 in ethanol is approximately 0.25 mg/ml and approximately 10 mg/ml in DMSO and DMF.

CU-115 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, CU-115 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. CU-115 has a solubility of approximately 0.15 mg/ml in a 1:5 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

CU-115 is a toll-like receptor 8 (TLR8) antagonist (IC $_{50}$ = 1.04 μ M). It is selective for TLR8 over TLR7 (IC $_{50}$ = >50 μ M). CU-115 inhibits increases in type I IFN transcriptional activity induced by the ssRNA nucleic acid ligands 3p-hpRNA or G3-YSD in a luciferase reporter assay when used at concentrations of 5 and 20 μM. It decreases TNF-α and IL-1β production induced by R-848 (Item No. 14806) in THP-1 cells when used at a concentration of $5 \mu M$.

Reference

1. Padilla-Salinas, R., Anderson, R., Sakaniwa, K., et al. Discovery of novel small molecule dual inhibitors targeting toll-like receptors 7 and 8. J. Med. Chem. 62(22), 10221-10244 (2019).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 10/15/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM