



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



Diazoxide-d₃

Item No. 29967

CAS Registry No.: 1432063-51-8

Formal Name: 7-chloro-3-(methyl-d₃)-4H-benzo[e]
[1,2,4]thiadiazine 1,1-dioxide

MF: C₈H₄ClD₃N₂O₂S

FW: 233.7

Chemical Purity: ≥95% (Diazoxide)


Deuterium

Incorporation: ≥99% deuterated forms (d₁-d₃); ≤1% d₀

Supplied as: A solid

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Diazoxide-d₃ is intended for use as an internal standard for the quantification of diazoxide (Item No. 14576) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Diazoxide-d₃ is supplied as a solid. A stock solution may be made by dissolving the diazoxide-d₃ in the solvent of choice, which should be purged with an inert gas. Diazoxide-d₃ is slightly soluble in DMSO.

Description

Diazoxide is an activator of sulfonylurea receptor 1 (SUR1) linked to ATP-sensitive potassium channel K_i_r6.2 (EC₅₀ = 14.1 μM in a FLIPR assay using HEK293 cells).¹ It also activates SUR2A/K_i_r6.2 and SUR2B/K_i_r6.2 channels in HEK293T cells in a patch-clamp assay when used at concentrations of 30 and 300 μM.² Diazoxide inhibits glucose-induced insulin release from isolated rat pancreatic β cells and induces relaxation of isolated rat aortic rings precontracted with potassium chloride (IC₅₀s = 22.6 and 22.4 μM, respectively).³ It reduces mean arterial pressure and cerebral blood flow in spontaneously hypertensive rats when administered intravenously as a 5 mg/kg bolus dose.⁴ Diazoxide (50 mg/kg, i.p.) increases blood glucose levels in mice.⁵ Formulations containing diazoxide have been used in the treatment of hypoglycemia.

References

1. Gopalakrishnan, M., Molinari, E.J., Char-Change, S., et al. *Br. J. Pharmacol.* **129**(7), 1323-1332 (2000).
2. Matsuoka, T., Matsushita, K., Katayama, Y., et al. *Circ. Res.* **87**(10), 873-880 (2000).
3. de Tullio, P., Becker, B., Boverie, S., et al. *J. Med. Chem.* **46**(15), 3342-3353 (2003).
4. Barry, D.I., Strandgaard, S., Graham, D.I., et al. *Eur. J. Clin. Invest.* **13**(3), 201-207 (1983).
5. Foy, J.M. and Furman, B.L. *Br. J. Pharmacol.* **47**(1), 124-132 (1973).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM