

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Methylophiopogonanone A

Item No. 30192

CAS Registry No.: 74805-92-8

Formal Name: (3R)-3-(1,3-benzodioxol-5-

> ylmethyl)-2,3-dihydro-5,7dihydroxy-6,8-dimethyl-4H-1-

benzopyran-4-one


MF: C₁₉H₁₈O₆ 342.3 FW: **Purity:** ≥98%

UV/Vis.: λ_{max} : 230 nm Supplied as: A crystalline solid

-20°C Storage: Stability: ≥4 years

Item Origin: Plant/Ophiopogonis radix

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Methylophiopogonanone A is supplied as a crystalline solid. A stock solution may be made by dissolving the methylophiopogonanone A in the solvent of choice, which should be purged with an inert gas. Methylophiopogonanone A is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of methylophiopogonanone A in ethanol is approximately 1 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Methylophiopogonanone A is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, methylophiopogonanone A should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Methylophiopogonanone A has a solubility of approximately 0.2 mg/ml in a 1:4 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Methylophiopogonanone A is a homoisoflavonoid that has been found in O. japonicus roots and has diverse biological activities.¹⁻⁴ It increases the uptake of rosuvastatin (Item Nos. 12029 | 18813) and atorvastatin (Item No. 10493) in HEK293T cells expressing human organic anion transporting polypeptide 1B1 (OATP1B1; EC₅₀s = 11.33 and 6 μ M, respectively).² Methylophiopogonanone A scavenges 2,2-diphenyl-1-picrylhydrazyl (DPPH; Item No. 14805) and ABTS (Item No. 27317) radicals in cell-free assays. In vivo, methylophiopogonanone A (10 mg/kg) decreases body weight gain and reduces serum and hepatic lipid levels in a rat model of high-fat diet-induced hyperlipidemia.³ It also reduces infarct size and myocardial apoptosis in mouse model of ischemia-reperfusion injury induced by transient occlusion of the left anterior descending coronary artery.4

References

- 1. Wang, Y., Liu, F., Liang, Z., et al. Iran J. Pharm. Res. 16(1), 357-365 (2017).
- 2. Chen, L., Liu, L., Chen, Y., et al. Xenobiotica 49(10), 1221-1228 (2019).
- 3. Li, Z., Wu, Y.Y., and Yu, B.X. Braz. J. Med. Biol. Res. 53(3), e9201 (2020).
- 4. He, F., Xu, B.L., Chen, C., et al. Acta. Pharmacol. Sin. 37(6), 763-771 (2016).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the mater can be found on our website.

Copyright Cayman Chemical Company, 12/12/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM