

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



**Domiphen** (bromide)

Item No. 30276

CAS Registry No.:	538-71-6
Formal Name:	N,N-dimethyl-N-(2-phenoxyethyl)-1-
	dodecanaminium, monobromide
Synonym:	NSC 39415
MF:	C <sub>22</sub> H <sub>40</sub> NO • Br
FW:	414.5
Purity:	≥98% •Br
UV/Vis.:	λ <sub>mav</sub> : 269 nm
Supplied as:	A crystalline solid
Storage:	-20°C
Stability:	≥4 years
Information represents	the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### Laboratory Procedures

Domiphen (bromide) is supplied as a crystalline solid. A stock solution may be made by dissolving the domiphen (bromide) in the solvent of choice, which should be purged with an inert gas. Domiphen (bromide) is soluble in organic solvents such as DMSO and methanol. It is also soluble in water. We do not recommend storing the aqueous solution for more than one day.

#### Description

Domiphen is a quaternary ammonium-containing cationic surfactant with antimicrobial activity.<sup>1,2</sup> It is active against A. viscosus, A. naeslundii, S. mutans, E. coli, and L. monocytogenes bacteria and inhibits C. neoformans yeast growth and spore germination.<sup>1-4</sup> It also inhibits the human-ether-a-go-go-related gene (hERG) channel (IC<sub>50</sub> = 1.5  $\mu$ M in a whole-cell patch-clamp assay).<sup>5</sup> Formulations containing domiphen have been used as antiseptics, disinfectants, and biocides in industrial, agricultural, veterinary, and clinical applications.

#### References

- 1. Baker, P.J., Coburn, R.A., Genco, R.J., et al. The in vitro inhibition of microbial growth and plague formation by surfactant drugs. J. Periodontal Res. 13(5), 474-485 (1978).
- 2. Müller, A., Rychli, K., Zaiser, A., et al. The Listeria monocytogenes transposon Tn6188 provides increased tolerance to various quaternary ammonium compounds and ethidium bromide. FEMS Microbiol. Lett. 361(2), 166-173 (2014).
- 3. Holdsworth, S.R. and Law, C.J. The major facilitator superfamily transporter MdtM contributes to the intrinsic resistance of Escherichia coli to quaternary ammonium compounds. J. Antimicrob. Chemother. 68(4), 831-839 (2013).
- 4. Ortiz, S.C., Huang, M., and Hull, C.M. Spore germination as a target for antifungal therapeutics. Antimicrob. Agents Chemother. 63(12), e00994-19 (2019).
- 5. Xia, M., Shahane, S.A., Huang, R., et al. Identification of quaternary ammonium compounds as potent inhibitors of hERG potassium channels. Toxicol. Appl. Pharmacol. 252(3), 250-258 (2011).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/24/2023

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM