

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Boc-NH-PEG₃ Acid

Item No. 30536

CAS Registry No.: 462100-06-7

Formal Name: 5,8,11-trioxa-2-azatridecanedioic

acid, 1-(1,1-dimethylethyl) ester

MF: $C_{13}H_{25}NO_{7}$ FW: 307.3 **Purity:** ≥95% Supplied as: A neat oil Storage: -20°C

Stability: ≥4 years Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

 $\mathsf{Boc}\text{-}\mathsf{NH}\text{-}\mathsf{PEG}_3$ acid is supplied as a neat oil. A stock solution may be made by dissolving the $\mathsf{Boc}\text{-}\mathsf{NH}\text{-}\mathsf{PEG}_3$ acid in the solvent of choice, which should be purged with an inert gas. Boc-NH-PEG₃ acid is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of Boc-NH-PEG₃ acid in these solvents is approximately 30, 1, and 5 mg/ml, respectively.

Description

 $\mathsf{Boc}\text{-}\mathsf{NH}\text{-}\mathsf{PEG}_3$ acid is a building $\mathsf{block}.^{1,2}$ It has been used as a linker in the synthesis of proteolysis-targeting chimeras (PROTACs) targeting MEK.¹ It has also been used as a spacer in the synthesis of radiolabeled probes for dual labeling of tumor integrin $\alpha V\beta 3$ and the gastrin-releasing peptide receptor (GRPR) in tumor tissue in vivo.2

References

- 1. Vollmer, S., Cunoosamy, D., Lv, H., et al. Design, synthesis, and biological evaluation of MEK PROTACs. J. Med. Chem. 63(1), 157-162 (2020).
- 2. Liu, Z., Yan, Y., Chin, F.T., et al. Dual integrin and gastrin-releasing peptide receptor targeted tumor imaging using ¹⁸F-labeled PEGylated RGD-bombesin heterodimer ¹⁸F-FB-PEG₃-Glu-RGD-BBN. J. Med. Chem. 52(2), 425-432 (2009).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 04/11/2025

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM