

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



MB-07811 Item No. 30670

CAS Registry No.: 852948-13-1

Formal Name: 4-[[4-[[(2R,4S)-4-(3-chlorophenyl)-

> 2-oxido-1,3,2-dioxaphosphorinan-2-yl]methoxy]-2,6-dimethylphenyl]

methyl]-2-(1-methylethyl)-phenol

≥4 years

Synonym: VK-2809 MF: $C_{28}H_{32}CIO_5P$ 515.0 FW: **Purity:** ≥98% Supplied as: A solid Storage: -20°C

HO.

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

MB-07811 is supplied as a solid. A stock solution may be made by dissolving the MB-07811 in the solvent of choice, which should be purged with an inert gas. MB-07811 is soluble in organic solvents such as acetonitrile and DMSO.

Description

Stability:

MB-07811 is a liver-targeted prodrug form of the thyroid hormone receptor β (TR β) agonist MB-07344. It is stable in extrahepatic tissues and converted to MB-07344 in the liver. MB-07811 binds to TRα1 and TR β 1 with K_i values of 12.5 and 14.6 μ M, respectively. Oral administration of MB-07811 reduces plasma cholesterol levels ($ED_{50} = 0.48 \text{ mg/kg}$) in a cholesterol-fed rat (CFR) model. It also reduces hepatic steatosis, as well as plasma levels of cholesterol, in Zucker diabetic fatty rats in a model of non-alcoholic fatty liver disease (NAFLD) when administered at a dose of 5 mg/kg per day. MB-07811 (10 mg/kg) reverses hepatic steatosis, reducing total hepatic lipids without decreasing body weight, in a mouse model of diet-induced obesity.

References

- 1. Boyer, S.H., Jiang, H., Jacintho, J.D., et al. Synthesis and biological evaluation of a series of liver-selective phosphonic acid thyroid hormone receptor agonists and their prodrugs. J. Med. Chem. 51(22), 7075-7093
- 2. Cable, E.E., Finn, P.D., Stebbins, J.W., et al. Reduction of hepatic steatosis in rats and mice after treatment with a liver-targeted thyroid hormone receptor agonist. Hepatology 49(2), 407-417 (2009).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 01/31/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM