



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)




PRODUCT INFORMATION



Fuziline

Item No. 30889

CAS Registry No.: 80665-72-1
Formal Name: (1 α ,6 α ,14 α ,15 α ,16 β)-20-ethyl-6,16-dimethoxy-4-(methoxymethyl)-aconitane-1,8,14,15-tetrol
Synonym: Senbusine C
MF: C₂₄H₃₉NO₇
FW: 453.6
Purity: $\geq$ 95%
Supplied as: A crystalline solid
Storage: -20°C
Stability: $\geq$ 2 years
Item Origin: Plant/*Radix Aconiti Carmichaeli*



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Fuziline is supplied as a crystalline solid. A stock solution may be made by dissolving the fuziline in the solvent of choice, which should be purged with an inert gas. Fuziline is soluble in chloroform and DMSO.

Description

Fuziline is a diterpene alkaloid that has been found in *A. lateral* and has cardioprotective activity.¹ It reduces sodium pentobarbital-induced cell death in primary neonatal rat cardiomyocytes when used at concentrations ranging from 0.1 to 10 μ M. Fuziline (0.5 μ M) reduces the production of reactive oxygen species (ROS) and apoptosis induced by isoproterenol (Item No. 15592) in H9c2 rat cardiomyocytes.² *In vivo*, fuziline (3 and 10 mg/kg) reduces isoproterenol-induced myocardial necrosis and fibrosis in rats.

References

1. Xiong, L., Peng, C., Xie, X.-F., et al. Alkaloids isolated from the lateral root of *Aconitum carmichaelii*. *Molecules* **17**(8), 9939-9946 (2012).
2. Fan, C.-L., Yao, Z.-H., Ye, M.-N., et al. Fuziline alleviates isoproterenol-induced myocardial injury by inhibiting ROS-triggered endoplasmic reticulum stress via PERK/eIF2 α /ATF4/Chop pathway. *J. Cell Mol. Med.* **24**(2), 1332-1344 (2020).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/18/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM