

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



McNA343

Item No. 30961

CAS Registry No.:	55-45-8	
Formal Name:	4-[[[(3-chlorophenyl)amino]carbonyl]	
	oxy]-N,N,N-trimethyl-2-butyn-1-	\sim
	aminium, monochloride	
MF:	$C_{14}H_{18}CIN_2O_2 \bullet CI$	
FW:	317.2	CI N O
Purity:	≥98%	N [×]
UV/Vis.:	λ _{max} : 239 nm	• Cl ⁻
Supplied as:	A solid	• 61
Storage:	-20°C	
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

McNA343 is supplied as a solid. A stock solution may be made by dissolving the McNA343 in the solvent of choice, which should be purged with an inert gas. McNA343 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of McNA343 in ethanol and DMF is approximately 15 mg/ml and approximately 25 mg/ml in DMSO.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of McNA343 can be prepared by directly dissolving the solid in aqueous buffers. The solubility of McNA343 in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

McNA343 is a partial agonist of muscarinic acetylcholine receptors (K_is = 5, 1, 5, 0.2, and 7.6 μ M for M₁₋₅ receptors, respectively, in radioligand binding assays)¹ It induces contraction of isolated human umbilical veins (EC₅₀ = 1.23 μ M), an effect that can be reversed by various M₁ muscarinic receptor selective antagonists. McNA343 also inhibits forskolin-induced cAMP production in CHO cells expressing M1 or M₄ receptors, as well as phosphatidylinositol hydrolysis in CHO-K1 cells expressing M₃ receptors and CHO cells expressing M₅ receptors. Intrathecal administration of McNA343 inhibits compulsive hindlimb neck-scratching behavior induced by 5'-guanidinonaltrindole (GNTI) in mice. It has nootropic activity, enhancing spontaneous working memory in the Y maze in wild-type mice, and it reverses learning and memory impairments induced by bulbectomy in mice in a passive avoidance test.

Reference

1. Mitchelson, F.J. The pharmacology of McN-A-343. Pharmacol. Ther. 135(2), 216-245 (2012).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/11/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM