



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION

## Clinofibrate

Item No. 31109

**CAS Registry No.:** 30299-08-2  
**Formal Name:** 2,2'-[cyclohexylidenebis(4,1-phenyleneoxy)]bis[2-methylbutanoic acid]  
**Synonym:** S-8527  
**MF:** C<sub>28</sub>H<sub>36</sub>O<sub>6</sub>  
**FW:** 468.6  
**Purity:** ≥95%  
**UV/Vis.:** λ<sub>max</sub>: 232 nm  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Clinofibrate is supplied as a crystalline solid. A stock solution may be made by dissolving the clinofibrate in the solvent of choice, which should be purged with an inert gas. Clinofibrate is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of clinofibrate in ethanol and DMF is approximately 30 mg/ml and approximately 15 mg/ml in DMSO.

Clinofibrate is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, clinofibrate should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Clinofibrate has a solubility of approximately 0.16 mg/ml in a 1:5 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

### Description

Clinofibrate is a hypolipidemic agent.<sup>1</sup> It inhibits HMG-CoA reductase (IC<sub>50</sub> = 0.47 mM) and stimulates the S-indan-1-ol dehydrogenase activity of the human liver 3α-hydroxysteroid dehydrogenase isoform AKR 1C4 in a concentration-dependent manner.<sup>2</sup> Clinofibrate (10 and 30 mg/kg) reduces serum cholesterol and triglyceride levels in rats fed a normal chow pellet diet or a semisynthetic diet containing sucrose as the only carbohydrate source.

### References

1. Toki, K., Nakamura, Y., Agatsuma, K., *et al.* Hypolipidemic action of a new aryloxy compound (S-8527) in rats. *Atherosclerosis* **18(1)**, 101-108 (1973).
2. Matsuura, K., Hara, A., Kato, M., *et al.* Activation of human liver 3α-hydroxysteroid dehydrogenase by clofibrate derivatives. *J. Pharmacol. Exp. Ther.* **285(3)**, 1096-1103 (1998).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/09/2020

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM