



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)




PRODUCT INFORMATION



CCG-215022

Item No. 31120

CAS Registry No.: 1813527-81-9
Formal Name: 4-[4-fluoro-3-[[[2-pyridinylmethyl]amino]carbonyl]phenyl]-1,2,3,4-tetrahydro-N-1H-indazol-5-yl-6-methyl-2-oxo-5-pyrimidinecarboxamide
MF: C₂₆H₂₂FN₇O₃
FW: 499.5
Purity: ≥98%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

CCG-215022 is supplied as a crystalline solid. A stock solution may be made by dissolving the CCG-215022 in the solvent of choice, which should be purged with an inert gas. CCG-215022 is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of CCG-215022 in these solvents is approximately 1 mg/ml.

CCG-215022 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, CCG-215022 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. CCG-215022 has a solubility of approximately 0.16 mg/ml in a 1:5 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

CCG-215022 is a G protein-coupled receptor kinase (GRK) inhibitor (IC₅₀s = 3.9, 0.15, and 0.38 μM for GRK1, GRK2, and GRK5, respectively).¹ It is selective for GRKs over protein kinase A (PKA; IC₅₀ = 120 μM). CCG-215022 (500 nM) increases contractility in isoproterenol-stimulated isolated mouse cardiomyocytes. It prevents the desensitization of histamine H₁ receptor- and purinergic P2Y₂ receptor-driven phospholipase C signaling in human ULTR myometrial cells and isolated rat mesenteric smooth muscle cells (MSMCs), respectively (IC₅₀s = 3.09 and 2.95 μM, respectively).²

References

1. RHoman, K.T., Waldschmidt, H.V., Glukhova, A., *et al.* Crystal structure of G protein-coupled receptor kinase 5 in complex with a rationally designed inhibitor. *J. Biol. Chem.* **290**(34), 20649-20659 (2015).
2. Rainbow, R.D., Brennan, S., Jackson, R., *et al.* Small-molecule G protein-coupled receptor kinase inhibitors attenuate G protein-coupled receptor kinase 2-mediated desensitization of vasoconstrictor-induced arterial contractions. *Mol. Pharmacol.* **94**(3), 1079-1091 (2018).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/16/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM