

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Macranthoidin B

Item No. 31533

CAS Registry No.: 136849-88-2 Formal Name: (3β,4α)-3-[(Ο-β-

> D-glucopyranosyl-(1→4)-O-β-Dglucopyranosyl- $(1\rightarrow 3)$ -O-6-deoxy- α -L-mannopyranosyl- $(1\rightarrow 2)-\alpha-L-$

> arabinopyranosyl) oxy]-23-hydroxyolean-12-en-28-oic acid, 6-O-β-Dglucopyranosyl-β-Dglucopyranosyl ester

Svnonvm: Macranthoiside I ${\sf C}_{65}{\sf H}_{106}{\sf O}_{32} \\ 1,399.5$ MF: FW:


Purity: ≥98%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Item Origin: Plant/Lonicera japonica

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Macranthoidin B is supplied as a crystalline solid. A stock solution may be made by dissolving the macranthoidin B in the solvent of choice, which should be purged with an inert gas. Macranthoidin B is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of macranthoidin B in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of macranthoidin B can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of macranthoidin B in PBS, pH 7.2, is approximately 5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Macranthoidin B is a triterpenoid saponin that has been found in Lonicerae and has anticancer activity.¹ It increases the production of reactive oxygen species (ROS) and induces apoptosis in HCT116 cells when used at concentrations ranging from 20 to 400 µM.

Reference

1. Fan, X., Rao, J., Zhang, Z., et al. Macranthoidin B modulates key metabolic pathways to enhance ROS generation and induce cytotoxicity and apoptosis in colorectal cancer. Cell Physiol. Biochem. 46(4), 1317-1330 (2018).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 09/03/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM