

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Prostaglandin $F_{2\alpha}$ - d_4

Item No. 316010

CAS Registry No.: 34210-11-2

Formal Name: 9α,11α,15S-trihydroxy-prosta-5Z,13E-

dien-1-oic-3,3,4,4-d₁ acid

Dinoprost-d₄, PGF_{2α}-d₄ Synonyms:

MF: C₂₀H₃₀D₄O₅ 358.5 FW: **Chemical Purity:** ≥98%


Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₄); \leq 1% d₀

Supplied as: A solution in methyl acetate

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Prostaglandin $F_{2\alpha}$ -d₄ (PGF_{2 α}-d₄) contains four deuterium atoms at the 3, 3', 4, and 4' positions. It is intended for use as an internal standard for the quantification of PGF_{2 α} (Item No. 16010) by GC- or LC-mass

 PGF_{2a} - d_4 is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of PGF20-d4 in these solvents is approximately 100 mg/ml.

 PGF_{2a} - d_4 is used as an internal standard for the quantification of PGF_{2a} by stable isotope dilution MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the weight indicated on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard PGF_{2a} by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Description

 $PGF_{2\alpha}$ is a widely distributed PG occurring in many species.¹⁻³ It causes contraction of vascular, bronchial, intestinal, and myometrial smooth muscle, and also exhibits potent luteolytic activity.¹ $PGF_{2\alpha}$ exherts its receptor mediated physiological activity at 50-100 nM.1 Maximal ovine myometrial contraction can be achieved at 125 nM PGF_{2a} in vitro.⁴

References

- 1. Samuelsson, B., Goldyne, M., Granström, E., et al. Annu. Rev. Biochem. 47, 997-1029 (1978).
- 2. Speroff, L. and Ramwell, P.W. Am. J. Obstet. Gynecol. 107, 1111-1130 (1970).
- 3. Watanabe, K., Iguchi, Y., Iguchi, S., et al. Proc. Natl. Acad. Sci. USA 83, 1583-1587 (1986).
- Crankshaw, D.J. and Gaspar, V. J. Reprod. Fertil. 103, 55-61 (1995).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/10/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM