

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Cilostazol-d₄ Item No. 31790

CAS Registry No.: 1215541-47-1

Formal Name: 6-(4-(1-(cyclohexyl-2,2,6,6-d₄)-

1H-tetrazol-5-yl)butoxy)-3,4-

dihydroquinolin-2(1H)-one

MF: $C_{20}H_{23}D_4N_5O_2$

373.5 FW:


Chemical Purity: ≥98% (Cilostazol)

Deuterium

 \geq 99% deuterated forms (d₁-d₄); \leq 1% d₀ Incorporation:

Supplied as: A solid -20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Cilostazol-d₄ is intended for use as an internal standard for the quantification of cilostazol (Item No. 15035) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Cilostazol-d₄ is supplied as a solid. A stock solution may be made by dissolving the cilostazol-d₄ in the solvent of choice, which should be purged with an inert gas. Cilostazol-d $_4$ is slightly soluble in DMSO and methanol.

Description

Cilostazol is a phosphodiesterase 3A (PDE3A) inhibitor ($IC_{50} = 0.2 \mu M$ for the platelet enzyme).¹ It is selective for PDE3A over PDE1, -2, -4, and 5 ($IC_{50}s = >5 \mu M$ for all). Cilostazol inhibits collagen- or ADP-induced aggregation of isolated rabbit platelets ($IC_{50}s = 29 \mu M$ and 31 μM , respectively). In vivo, cilostazol (30 mg/kg) reduces thrombus formation by 84% in a mouse model of pulmonary thromboembolism. It reduces cardiac fibrosis and prevents the development of diastolic dysfunction and cardiac hypertrophy induced by a high-fat diet and angiotensin II in mice. Formulations containing cilostazol have been used in the treatment of the symptoms of intermittent claudication in peripheral vascular disease.

References

- 1. Schrör, K. The pharmacology of cilostazol. Diabetes Obes. Metab. 4 (Suppl 2), S14-S19 (2002).
- 2. Koga, Y., Kihara, Y., Okada, M., et al. 2(1H)-Quinolinone derivatives as novel anti-arteriostenotic agents showing anti-thrombotic and anti-hyperplastic activities. Bioorg. Med. Chem. Lett. 8(12), 1471-1476 (1998).
- 3. Reddy, S.S., Agarwal, H., and Barthwal, M.K. Cilostazol ameliorates heart failure with preserved ejection fraction and diastolic dysfunction in obese and non-obese hypertensive mice. J. Mol. Cell. Cardiol. 123, 46-57 (2018).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 09/30/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM