

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



GW 791343

Item No. 32882

CAS Registry No.: 1019779-04-4

Formal Name: 2-[(3,4-difluorophenyl)amino]-N-[2-methyl-

5-(1-piperazinylmethyl)phenyl]-acetamide,


dihydrochloride

MF: C₂₀H₂₄F₂N₄O • 2HCl

FW: 447.4 **Purity:** ≥98% UV/Vis.: λ_{max} : 235 nm Supplied as: A crystalline solid

Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

GW 791343 is supplied as a crystalline solid. A stock solution may be made by dissolving the GW 791343 in the solvent of choice, which should be purged with an inert gas. GW 791343 is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of GW 791343 in these solvents is approximately 1 and 10 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of GW 791343 can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of GW 791343 in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

GW 791343 is an allosteric modulator of the purinergic P2X₇ receptor.¹ It acts as a negative allosteric modulator at human P2X₇ receptors but as a positive allosteric modulator at rat P2X₇ receptors. GW 791343 (10 and 30 μM) reduces ethidium accumulation induced by the P2X receptor agonists ATP (Item No. 14498) or BzATP (Item No. 15577) in HEK293 cells expressing the human receptor. In contrast, it increases ethidium accumulation induced by ATP or BzATP in HEK293 cells expressing the rat receptor when used at the same concentrations. GW 791343 (5 μ M) potentiates the circadian release of ATP in rat organotypic suprachiasmatic nucleus (SCN) slice cultures.²

References

- 1. Michel, A.D., Chambers, L.J., and Walter, D.S. Negative and positive allosteric modulators of the P2X₇ receptor. Br. J. Pharmacol. 153(4), 737-750 (2008).
- Svobodova, I., Bhattaracharya, A., Ivetic, M., et al. Circadian ATP Release in Organotypic Cultures of the Rat Suprachiasmatic Nucleus Is Dependent on P2X₇ and P2Y Receptors. Front. Pharmacol. 9, 192 (2018).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 10/18/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM