



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



NADPH Cytochrome P450 Reductase (human, recombinant)


Item No. 33015

Overview and Properties


Synonyms: CPR, CYPOR, NADPH Hemoprotein Reductase, NCPR, P450R, POR
Source: Active recombinant human C-terminal His-tagged NADPH cytochrome P450 reductase expressed in *E. coli*
Amino Acids: 24-677
Uniprot No.: P16435
Molecular Weight: 76.4 kDa
Storage: -80°C (as supplied)
Stability: ≥1 year
Purity: ≥90% estimated by SDS-PAGE
Supplied in: 50 mM potassium phosphate, pH 7.6, with 10% glycerol
Protein Concentration: *batch specific* mg/ml
Activity: *batch specific* U/ml
Specific Activity: *batch specific* U/mg
Unit Definition: One unit is defined as the amount of enzyme required to reduce 1 nmol of cytochrome c per minute at 25°C in 25 mM potassium phosphate, pH 7.6.

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Images



Lane 1: MW Markers
Lane 2: NADPH Cytochrome P450 Reductase (human, recombinant) (2 µg)
Lane 3: NADPH Cytochrome P450 Reductase (human, recombinant) (4 µg)



Typical UV-visible absorption spectra of purified NADPH Cytochrome P450 Reductase (human, recombinant). The presence of FAD loaded domains in NADPH Cytochrome P450 Reductase (human, recombinant) are verified spectrophotometrically by FAD absorption at 463 nm using an extinction coefficient of $0.0136 \mu\text{M}^{-1} \text{cm}^{-1}$.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.


SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/25/2023

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



NADPH Cytochrome P450 Reductase (human, recombinant) activity was determined through the reduction of cytochrome *c* from bovine heart (Sigma-Aldrich, Product No. C3131) using β -NADPH (Sigma-Aldrich, Product No. N7505) as the cofactor. The reduction of cytochrome *c* results in the increase in absorbance at 550 nm, thereby allowing activity to be monitored.

Description

NADPH cytochrome P450 reductase (POR) is a flavoprotein and key electron donor to cytochrome P450s (CYPs).^{1,2} It is composed of an N-terminal membrane anchor domain and FMN-binding domain, which is linked to a C-terminal FAD-binding domain and NADPH-binding domain and localized to the cytosolic side of the endoplasmic reticulum.^{1,3,4} POR transfers electrons from NADPH through FAD and FMN coenzymes into the iron of the prosthetic heme group of CYPs.^{1,3} Cortical levels of POR are increased in the APPswe/PSEN1dE9 transgenic mouse model of Alzheimer's disease.⁵ Deficiency in POR is associated with ambiguous genitalia and the disruption of steroid metabolism in patients with Antley-Bixler syndrome.⁶ Cayman's NADPH Cytochrome P450 Reductase (human, recombinant) can be used for enzyme activity assays.

References

1. Laursen, T., Jensen, K., and Møller, B.L. Conformational changes of the NADPH-dependent cytochrome P450 reductase in the course of electron transfer to cytochromes P450. *Biochim. Biophys. Acta.* **1814(1)**, 132-138 (2011).
2. Backes, W.L. and Kelley, R.W. Organization of multiple cytochrome P450s with NADPH-cytochrome P450 reductase in membranes. *Pharmacol. Ther.* **98(2)**, 221-233 (2003).
3. Jensen, K. and Møller, B.L. Plant NADPH-cytochrome P450 oxidoreductases. *Phytochemistry* **71(2-3)**, 132-141 (2010).
4. Barnaba, C., Gentry, K., Sumangala, N., *et al.* The catalytic function of cytochrome P450 is entwined with its membrane-bound nature. *F1000Res.* **6(F1000 Faculty Rev)**, 662 (2017).
5. Yao, Y., Huang, J.-Z., Chen, L., *et al.* *In vivo* and *in vitro* studies on the roles of p38 mitogen-activated protein kinase and NADPH-cytochrome P450 reductase in Alzheimer's disease. *Exp. Ther. Med.* **14(5)**, 4755-4760 (2017).
6. Pandey, A.V. and Flück, C.E. NADPH P450 oxidoreductase: Structure, function, and pathology of diseases. *Pharmacol. Ther.* **138(2)**, 229-254 (2013).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM