



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



MF438

Item No. 33349

CAS Registry No.: 921605-87-0
Formal Name: 3-(5-methyl-1,3,4-thiadiazol-2-yl)-6-[4-[2-(trifluoromethyl)phenoxy]-1-piperidinyl]-pyridazine
MF: C₁₉H₁₈F₃N₅OS
FW: 421.4
Purity: ≥98%
UV/Vis.: λ_{max}: 224, 316 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

MF438 is supplied as a crystalline solid. A stock solution may be made by dissolving the MF438 in the solvent of choice, which should be purged with an inert gas. MF438 is soluble in the organic solvent dimethyl formamide at a concentration of approximately 30 mg/ml.

Description

MF438 is an inhibitor of stearoyl-CoA desaturase 1 (SCD1; IC₅₀ = 2.3 nM for the rat enzyme), an enzyme that catalyzes the formation of a *cis* double bond at the Δ⁹ position of stearoyl-CoA to produce oleoyl-CoA.^{1,2} It reduces spheroid formation in NCI H460 and patient-derived non-small cell lung cancer (NSCLC) cells in a concentration-dependent manner, an effect that can be blocked by oleic acid (Item Nos. 90260 | 24659).³ MF438 (1 μM) decreases the activity of aldehyde dehydrogenase 1A1 (ALDH1A1), a marker of cancer stem cells (CSCs), in NCI H460-derived spheroids when used at a concentration of 1 μM. It acts synergistically with cisplatin (Item No. 13119) to induce apoptosis and autophagy in patient-derived NSCLC cells.⁴

References

1. Léger, S., Black, C., Deschenes, D., *et al.* Synthesis and biological activity of a potent and orally bioavailable SCD inhibitor (MF-438). *Bioorg. Med. Chem. Lett.* **20**(2), 499-502 (2010).
2. Isabel, E., Powell, D.A., Black, W.C., *et al.* Biological activity and preclinical efficacy of azetidiny pyridazines as potent systemically-distributed stearoyl-CoA desaturase inhibitors. *Bioorg. Med. Chem. Lett.* **21**(1), 479-483 (2011).
3. Noto, A., Raffa, S., De Vitis, C., *et al.* Stearoyl-CoA desaturase-1 is a key factor for lung cancer-initiating cells. *Cell Death Dis.* **4**(12), e947 (2013).
4. Pisanu, M.E., Noto, A., De Vitis, C., *et al.* Blockade of stearoyl-CoA-desaturase 1 activity reverts resistance to cisplatin in lung cancer stem cells. *Cancer Lett.* **406**, 93-104 (2017).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/04/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM