



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)




PRODUCT INFORMATION



Batabulin (sodium salt)

Item No. 33487

CAS Registry No.: 195533-98-3
Formal Name: 2,3,4,5,6-pentafluoro-N-(3-fluoro-4-methoxyphenyl)-benzenesulfonamide, monosodium salt
Synonym: T138067
MF: C₁₃H₆F₆NO₃S • Na
FW: 393.2
Purity: ≥98%
UV/Vis.: λ_{max}: 235 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Batabulin (sodium salt) is supplied as a crystalline solid. A stock solution may be made by dissolving the batabulin (sodium salt) in the solvent of choice, which should be purged with an inert gas. Batabulin (sodium salt) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of batabulin (sodium salt) in these solvents is approximately 30 mg/ml.

Batabulin (sodium salt) is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, batabulin (sodium salt) should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Batabulin (sodium salt) has a solubility of approximately 0.25 mg/ml in a 1:3 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Batabulin is an inhibitor of tubulin polymerization.¹ It covalently binds to cysteine 239 (Cys²³⁹) of the β1, β2, and β4 tubulin isotypes and inhibits polymerization of bovine brain tubulin (IC₅₀ = 2 μM) in cell-free assays. Batabulin (30, 100, and 300 nM) induces cytoskeletal collapse and cell cycle arrest at the G₂/M phase in MCF-7 breast cancer cells. It is cytotoxic against a panel of drug-sensitive and -resistant cancer cell lines (IC₅₀s = 11-165 nM). Batabulin (40 mg/kg) reduces tumor volume in CCRF CEM lymphoblastic leukemia and CEM/VBL100 multidrug-resistant lymphoblastic leukemia mouse xenograft models.

Reference

1. Shan, B., Medina, J.C., Santha, E., *et al.* Selective, covalent modification of β-tubulin residue Cys-239 by T138067, an antitumor agent with *in vivo* efficacy against multidrug-resistant tumors. *Proc. Natl. Acad. Sci. USA* **96**(10), 5686-5691 (1999).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/29/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 • USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM