

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Thromboxane A Synthase (human, recombinant)

Item No. 33678

Overview and Properties

Synonyms: CYP5A1, Cytochrome P450 5A1, TBXAS1, TXAS, TXA Synthase

Source: Active recombinant C-terminal His-tagged thromboxane A synthase expressed in E. coli

Amino Acids: 30-533 (full length) with a modified N-terminus and F-G loop

Uniprot No.: P24557 Molecular Weight: 59.1 kDa

-80°C (as supplied) Storage:

Stability: ≥6 months

Purity: batch specific (≥90% estimated by SDS-PAGE)

Supplied in: 50 mM Tris, pH 7.4, with 5% glycerol, 0.2 mM DTT, and 0.2 mM EDTA


Protein

Concentration: batch specific mg/ml

Activity: Confirmed

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Images



Lane 2: Thromboxane A Synthase (2 µg) Lane 3: Thromboxane A Synthase (4 µg)

SDS-PAGE Analysis of Thromboxane A Synthase.



Panel A: Assay using 18 µM TXAS

Lane 1: PGH,

Lane 2: PGH₂ conversion for 30 min. Lane 3: PGH₂ conversion for one hour.

Lane 4: 12(S)-HHTrE (Item No. 34590)

Lane 5: TXB₂ (Item No. 19030)



Panel B: Assay using 9 µM TXAS

Lane 1: PGH,

Lane 2: NO substrate control Lane 3: PGH₂ conversion for 30 min. Lane 4: PGH₂ conversion for one hour.

Lane 5: TXB₂ (Item No. 19030)

Thin layer chromatographic analysis of Prostaglandin H₂ (PGH₂; Item No. 17020) conversion by recombinant TXAS (human, recombinant). In vitro conversion of PGH, (50 μ M) was done using 18 μ M (Panel A) and 9 μ M (Panel B) in 250 μ L of assay (30 μm) was done using 16 μm (raner A) and 9 μm (raner b) in 250 μc of assay buffer (10 mM potassium phosphate buffer, pH 7.4, with 0.2 mM DTT, 0.5 mM EDTA and 10% glycerol). The concentration of the protein was based on the Ferrous-CO complexed form of the protein. Analysis of the product was performed in a TLC plate using solvent 40:60:1 ethyl acetate:heptane:acetic acid at 4°C. The plates were visualized by spraying 5% sulfuric acid in methanol.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/20/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Thromboxane A synthase (TXAS), also known as cytochrome P450 (CYP) isoform CYP5A1, is the enzyme that catalyzes the isomerization of prostaglandin H_2 (PGH $_2$; Item No. 17020) into thromboxane A_2 (TXA $_2$), a vasoconstrictor and an inducer of platelet aggregation. It also catalyzes the cleavage of PGH $_2$ into malondialdehyde (MDA) and 12(S)-hydroxyheptadecatrienoic acid (12(S)-HHT; Item No. 34590), a leukotriene B_4 (LTB $_4$) receptor 2 (BLT $_2$) agonist that also has a role in platelet aggregation. TXAS exists as a monomer and is composed of an N-terminal membrane anchor domain, a heme-binding catalytic residue, and several substrate-binding residues. It is expressed in numerous cells, including platelets, monocytes, and macrophages, as well as several tissues, and is localized to the endoplasmic reticulum. TXAS undergoes suicide inactivation during catalysis. Mice deficient in TXAS exhibit prolonged bleeding time and defective platelet aggregation. Cayman's Thromboxane A Synthase (human, recombinant) protein can be used for enzyme activity assay and Western blot (WB) applications. To construct this protein, the N-terminal amino acids 1-29 of human CYP5A1 were removed and replaced with a hydrophilic sequence. The F-G loop of TXAS was modified using sequences from rabbit CYP2C5 and CYP2C3. The C-terminal of this protein contains 6x-His tag followed by a stop codon.

References

- 1. Okuno, T. and Yokomizo, T. Biological functions of 12(S)-hydroxyheptadecatrienoic acid as a ligand of leukotriene B_4 receptor 2. *Inflamm. Regen.* **38**, 29 (2018).
- 2. Hajeyah, A.A., Griffiths, W.J., Wang, Y., et al. The biosynthesis of enzymatically oxidized lipids. Front. Endocrinol. (Lausanne) 11, 591819 (2020).
- 3. Ershov, P.V., Yablokov, E., Zgoda, V., et al. A new insight into subinteractomes of functional antagonists: Thromboxane (CYP5A1) and prostacyclin (CYP8A1) synthases. *Cell Biol. Int.* **45(6)**, 1175-1182 (2021).
- 4. Ruan, K.H., Milfield, K., Kulmacz, R.J., *et al.* Comparison of the construction of a 3-D model for human thromboxane synthase using P450cam and BM-3 as templates: Implications for the substrate binding pocket. *Protein Eng.* **7(11)**, 1345-1351 (1994).
- 5. Seo, M.-J. and Oh, D.-K. Prostaglandin synthases: Molecular characterization and involvement in prostaglandin biosynthesis. *Prog. Lipid Res.* **66**, 50-68 (2017).
- 6. Yu, I.-S., Lin, S.-R., Huang, C.-C., et al. TXAS-deleted mice exhibit normal thrombopoiesis, defective hemostasis, and resistance to arachidonate-induced death. *Blood* **104(1)**, 135-142 (2004).

PHONE: [800] 364-9897 [734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM