



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION

rac-Desethyl Oxybutynin (hydrochloride)

Item No. 34262

CAS Registry No.: 81039-77-2

Formal Name: *α*-cyclohexyl-*α*-hydroxy-benzeneacetic acid, 4-(ethylamino)-2-butyn-1-yl ester, monohydrochloride

Synonym: (R,S)-DEOB

MF: C₂₀H₂₇NO₃ • HCl


FW: 365.9

Purity: ≥95%

Supplied as: A solid

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

rac-Desethyl oxybutynin (hydrochloride) is supplied as a solid. A stock solution may be made by dissolving the *rac*-desethyl oxybutynin (hydrochloride) in the solvent of choice, which should be purged with an inert gas. *rac*-Desethyl oxybutynin (hydrochloride) is slightly soluble in methanol and chloroform.

Description

rac-Desethyl oxybutynin is an active metabolite of the muscarinic acetylcholine receptor (mAChR) antagonist oxybutynin (Item No. 23825).¹⁻³ *rac*-Desethyl oxybutynin is formed from oxybutynin by the cytochrome P450 (CYP) isoform CYP3A4.² It binds to mAChRs in isolated rat bladder, submaxillary gland, heart, and colon with K_i values of 5.38, 3.12, 16, and 5.59 nM, respectively. *rac*-Desethyl oxybutynin (0.4 and 0.8 mg/kg) reduces micturition pressure in rats.³

References

1. Lindeke, B., Hallström, G., Johansson, C., et al. Metabolism of oxybutynin: Establishment of desethyloxybutynin and oxybutynin N-oxide formation in rat liver preparation using deuterium substitution and gas chromatographic mass spectrometric analysis. *Biomed. Mass Spectrom.* **8**(10), 506-513 (1981).
2. Oki, T., Kawashima, A., Uchida, M., et al. In vivo demonstration of muscarinic receptor binding activity of N-desethyl-oxybutynin, active metabolite of oxybutynin. *Life Sci.* **76**(21), 2445-2456 (2005).
3. Mizushima, H., Kinoshita, K., Abe, K., et al. Pharmacokinetics/pharmacodynamics analysis of the relationship between the *in vivo* micturition pressure and receptor occupancy of (R)-oxybutynin and its metabolite in rats. *Biol. Pharm. Bull.* **30**(5), 955-962 (2007).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.