

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



8(R)-HETE

Item No. 34350

CAS Registry No.: 105500-09-2

Formal Name: 8R-hydroxy-5Z,9E,11Z,14Z-

eicosatetraenoic acid

MF: $C_{20}H_{32}O_3$ FW: 320.5 **Purity:** ≥98%

Stability: ≥1 year at -20°C Supplied as: A solution in ethanol Special Conditions: Oxygen and light sensitive

Laboratory Procedures

For long term storage, we suggest that 8(R)-HETE be stored as supplied at -20°C. It should be stable for at least one year.

8(R)-HETE is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of 8(R)-HETE in these solvents is approximately 50 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 8(R)-HETE is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of 8(R)-HETE in PBS (pH 7.2) is approximately 1 mg/ml. Store aqueous solutions of 8(R)-HETE on ice and use within 12 hours of preparation.

Description

8(R)-HETE is a metabolite of arachidonic acid by the 8-lipoxygenase pathway in marine organisms.^{1,2} It induces maturation of starfish oocytes at a concentration of 0.071 µM. This activity if neglible in the 8(S)-enantiomer. The precursor of 8(R)-HETE, 8(R)-HPETE, is the putative intermediate in the biosynthesis of marine prostaglandins via the allene oxide synthase pathway.² Stereochemical assignment of the (R) enantiomer is based on comparison of chiral HPLC retention times to published results.³

References

- 1. Meijer, L., Brash, A.R., Bryant, R.W., et al. Stereospecific induction of starfish oocyte maturation by (8R)-hydroxyeicosatetraenoic acid. J. Biol. Chem. 261, 17040-17047 (1986).
- 2. Brash, A.R., Baertschi, S.W., Ingram, C.D., et al. On non-cyclooxygenase prostaglandin synthesis in the sea whip coral, Plexaura Homomalla: an 8(R)-lipoxygenase pathway leads to formation of an α -ketol and a racemic prostanoid. J. Biol. Chem. 262, 15829-15839 (1987).
- 3. Schneider, C., Yu, Z., Boeglin, W.E., et al. Enantiomeric separation of hydroxy and hydroperoxy eicosanoids by chiral column chromatography. Method. Enzymol. 433, 145-157 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/21/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM