



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Saikosaponin C

Item No. 34599

CAS Registry No.: 20736-08-7
Formal Name: (3 β ,16 β)-13,28-epoxy-16-hydroxyolean-11-en-3-yl O-6-deoxy- α -L-mannopyranosyl-(1 \rightarrow 4)-O-[β -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranoside

MF: C₄₈H₇₈O₁₇

FW: 927.1


Purity: \geq 98%

Supplied as: A solid

Storage: -20°C

Stability: \geq 2 years

Item Origin: Plant/Radix Bupleuri



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Saikosaponin C is supplied as a solid. A stock solution may be made by dissolving the saikosaponin C in the solvent of choice, which should be purged with an inert gas. Saikosaponin C is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of saikosaponin C in ethanol is approximately 1 mg/ml and approximately 10 mg/ml in DMSO and DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of saikosaponin C can be prepared by directly dissolving the solid in aqueous buffers. The solubility of saikosaponin C in PBS (pH 7.2) is approximately 1 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Saikosaponin C is a saikosaponin that has been found in *B. falactum* and has anti-inflammatory and analgesic activities.^{1,2} It reduces basal production of prostaglandin E₂ (PGE₂; Item No. 14010) in isolated rat peritoneal macrophages when used at a concentration of 100 μ g/ml.¹ Saikosaponin C (5 mg/kg per day) decreases acetic acid-induced writhing in repeated cold stressed (SART) mice.²

References

- Ohuchi, K., Watanabe, M., Ozeki, T., *et al.* Pharmacological influence of saikosaponins on prostaglandin E₂ production by peritoneal macrophages. *Planta Med.* **51**(3), 208-212 (1985).
- Kita, T., Hata, T., Itoh, E., *et al.* Analgesic and other pharmacologic actions of saiko-saponin in repeated cold stressed (SART stressed) animals. *J. Pharmacobiodyn.* **3**(6), 269-280 (1980).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/04/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM