



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION

Norfluoxetine-d₅ (hydrochloride)

Item No. 34742

CAS Registry No.: 1185132-92-6

Formal Name: γ-[4-(trifluoromethyl)phenoxy]-benzenepropan-1,1,2,2,3-d₅-amine, monohydrochloride

Synonym: Desmethylfluoxetine-d₅

MF: C₁₆H₁₁D₅F₃NO • HCl

FW: 336.8

Chemical Purity: ≥98% (Norfluoxetine)


Deuterium

Incorporation: ≥99% deuterated forms (d₁-d₅); ≤1% d₀

Supplied as: A solid

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Norfluoxetine-d₅ (hydrochloride) is intended for use as an internal standard for the quantification of norfluoxetine (Item No. 15900) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Norfluoxetine-d₅ (hydrochloride) is supplied as a solid. A stock solution may be made by dissolving the norfluoxetine-d₅ (hydrochloride) in the solvent of choice, which should be purged with an inert gas. Norfluoxetine-d₅ (hydrochloride) is slightly soluble in chloroform and DMSO.

Description

Norfluoxetine is an active metabolite of the antidepressant fluoxetine.¹ It is formed from fluoxetine by the cytochrome P450 (CYP) isoforms CYP2C9, CYP2C19, and CYP3A.² Norfluoxetine inhibits serotonin (5-HT) uptake in rat brain synaptosomal membrane preparations (K_i = 44.7 nM) and isolated human platelets (IC₅₀ = ~15 nM).¹ It has been found in the tissues of fish exposed to wastewater effluent.³

References

- Wong, D.T., Bymaster, F.P., Reid, L.R., et al. Norfluoxetine enantiomers as inhibitors of serotonin uptake in rat brain. *Neuropsychopharmacology* 8(4), 337-344 (1993).
- Hiemke, C. and Härtter, S. Pharmacokinetics of selective serotonin reuptake inhibitors. *Pharmacol. Ther.* 85(1), 11-28 (2000).
- Brooks, B.W., Chambliss, C.K., Stanley, J.K., et al. Determination of select antidepressants in fish from an effluent-dominated stream. *Environ. Toxicol. Chem.* 24(2), 464-469 (2005).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.