

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



L-Methionine-d₂

Item No. 34826

CAS Registry No.: 13010-53-2

Formal Name: S-(methyl-d₃)-L-homocysteine

Synonym: (S)-Methionine-d₂ MF: $C_5H_8D_3NO_2S$


FW: 152.2 **Chemical Purity:** ≥95% (L-methionine)

Deuterium

Incorporation: ≥99% deuterated forms (d_1-d_3) ; ≤1% d_0

UV/Vis.: λ_{max} : 570 nm Supplied as: A solid -20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

L-Methionine-d₃ is intended for use as an internal standard for the quantification of L-methionine by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Description

L-Methionine is an essential amino acid.^{1,2} It is catabolized by methionine adenosyltransferase 2A (MAT2A), yielding the universal methyl donor S-(5'-adenosyl-L-methionine (SAM; Item Nos. 16376 | 17595).² L-methionine has roles in purine and pyrimidine synthesis, polyamine synthesis, and glutathione production, as well as in various metabolic diseases and aging. Dietary methionine restriction (MR) prevents accelerated aging in a mouse model of progeria and increases the lifespan of yeast, Drosophila, C. elegans, mice, and rats. 1,2 MR also improves insulin sensitivity and reduces adiposity in mice, as well as inhibits the growth and proliferation of several cancer cell lines.

References

- 1. Wanders, D., Hobson, K., and Ji, X. Methionine restriction and cancer biology. Nutrients 12(3), 684 (2020).
- 2. Sanderson, S.M., Gao, X., Dai, Z., et al. Methionine metabolism in health and cancer: A nexus of diet and precision medicine. Nat. Rev. Cancer 19(11), 625-637 (2019).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/20/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM