



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)




PRODUCT INFORMATION



L-Leucine-d₁₀ Item No. 34843

CAS Registry No.: 106972-44-5
Formal Name: L-leucine-2,3,3,4,5,5,5,5',5',5'-d₁₀
Synonyms: Leu-d₁₀, Pentanoic Acid-d₁₀
MF: C₆H₃D₁₀NO₂
FW: 141.2
Chemical Purity: ≥98% (L-Leucine)
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₁₀); ≤1% d₀
Supplied as: A solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

L-Leucine-d₁₀ is intended for use as an internal standard for the quantification of L-leucine (Item No. 34342) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Description

L-Leucine is an essential amino acid.¹ It increases basal insulin secretion and decreases glucose-induced insulin release in INS-1E rat insulinoma cells when used at concentrations of 1, 5, and 10 mM.² L-Leucine (1, 5, and 10 mM) decreases triglyceride levels and increases cholesterol accumulation in INS-1E cells. It stimulates skeletal muscle protein synthesis in exercised rats, as well as in food-deprived rats in an mTOR-dependent manner.^{3,4} Formulations containing L-leucine have been used as dietary supplements.

References

1. Duan, Y., Li, F., Li, Y., *et al.* The role of leucine and its metabolites in protein and energy metabolism. *Amino Acids* **48**(1), 41-51 (2016).
2. Liu, Z., Jeppesen, P.B., Gregersen, S., *et al.* Chronic exposure to leucine *in vitro* induces β-cell dysfunction in INS-1E cells and mouse islets. *J. Endocrinol.* **215**(1), 79-88 (2012).
3. Anthony, J.C., Yoshizawa, F., Anthony, T.G., *et al.* Leucine stimulates translation initiation in skeletal muscle of postabsorptive rats via a rapamycin-sensitive pathway. *J. Nutr.* **130**(10), 2413-2419 (2000).
4. Anthony, J.C., Anthony, T.G., and Layman, D.K. Leucine supplementation enhances skeletal muscle recovery in rats following exercise. *J. Nutr.* **129**(6), 1102-1106 (1999).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/08/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM