



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



IDO-IN-2

Item No. 34876

CAS Registry No.: 1668565-74-9
Formal Name: N-[4-[bis(2-methylpropyl)amino]-2'-(2H-tetrazol-5-yl)[1,1'-biphenyl]-3-yl]-N'-(4-methylphenyl)-urea
Synonym: PCC0208009
MF: C₂₉H₃₅N₇O
FW: 497.6
Purity: ≥98%
UV/Vis.: λ_{max}: 257 nm
Supplied as: A solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

IDO-IN-2 is supplied as a solid. A stock solution may be made by dissolving the IDO-IN-2 in the solvent of choice, which should be purged with an inert gas. IDO-IN-2 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of IDO-IN-2 in ethanol is approximately 2 mg/ml and approximately 10 mg/ml in DMSO and DMF.

Description

IDO-IN-2 is an inhibitor of indoleamine 2,3-dioxygenase 1 (IDO1; IC₅₀ = 0.097 μM in HeLa cells).¹ *In vivo*, IDO-IN-2 (100 mg/kg) decreases intratumor levels of Ki67, a marker of cell proliferation, and reduces tumor weight in a GL261 murine glioma heterotopic transplantation model.² It also decreases mechanical and thermal hypersensitivity, improves novel object recognition, and decreases anterior cingulate cortex (ACC) and amygdala levels of IDO1 in a rat model of neuropathic pain induced by spinal nerve ligation (SNL).³

References

1. Williams, D.K., Markwalder, J.A., Balog, A.J., *et al.* Development of a series of novel *o*-phenylenediamine-based indoleamine 2,3-dioxygenase 1 (IDO1) inhibitors. *Bioorg. Med. Chem. Lett.* **28**(4), 732-736 (2018).
2. Sun, S., Du, G., Xue, J., *et al.* PCC0208009 enhances the anti-tumor effects of temozolomide through direct inhibition and transcriptional regulation of indoleamine 2,3-dioxygenase in glioma models. *Int. J. Immunopathol. Pharmacol.* **32**, 2058738418787991 (2018).
3. Wang, Y., Li, C.-M., Han, R., *et al.* PCC0208009, an indirect IDO1 inhibitor, alleviates neuropathic pain and co-morbidities by regulating synaptic plasticity of ACC and amygdala. *Biochem. Pharmacol.* **177**, 113926 (2020).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/20/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM