

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## Dimidium (bromide)

Item No. 35816

CAS Registry No.: 518-67-2

Formal Name: 3,8-diamino-5-methyl-6-phenyl-

phenanthridinium, monobromide

Synonym: Phenanthridinium Compound 1553


MF:  $C_{20}H_{18}N_3 \bullet Br$ 

380.3 FW: ≥98% **Purity:** 

UV/Vis.:  $\lambda_{max}$ : 214, 295 nm

Supplied as: A solid Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



#### **Laboratory Procedures**

Dimidium (bromide) is supplied as a solid. A stock solution may be made by dissolving the dimidium (bromide) in the solvent of choice, which should be purged with an inert gas. Dimidium (bromide) is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of dimidium (bromide) in these solvents is approximately 3 and 2 mg/ml, respectively.

#### Description

Dimidium is a trypanocide. <sup>1,2</sup> In vivo, dimidium (0.25-0.7 mg/kg) reduces parasite burden in T. congolenseinfected mice. It also completely clears trypanosomes from the peripheral blood in zebu oxen naturally infected with T. congolense when administered intramuscularly at a dose of 0.75 mg/kg.<sup>2</sup> Formulations containing dimidium have previously been used in the treatment of trypanosomiasis in cattle.

### References

- 1. Woolfe, G. The trypanocidal action of phenanthridine compounds. Ann. Trop. Med. Parasitol. 46(4), 285-288 (1952).
- 2. Wilde, J.K.H. Experimental use of dimidium bromide (1553) in the treatment of T. congolense infection in the zebu cattle. J. Comp. Pathol. Ther. 59(1), 48-53 (1949).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

## WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/13/2022

#### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM