



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



Timosaponin AI

Item No. 36174

CAS Registry No.: 68422-00-4

Formal Name: (3 β ,5 β ,25S)-spirostan-3-yl β -D-galactopyranoside

Synonym: Sarsasapogenin-3-O- β -galactopyranoside

MF: C₃₃H₅₄O₈

FW: 578.8


Purity: \geq 98%

Supplied as: A solid

Storage: -20°C

Stability: \geq 4 years

Item Origin: Plant/*Anemarrhena asphodeloides*



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Timosaponin AI is supplied as a solid. A stock solution may be made by dissolving the timosaponin AI in the solvent of choice, which should be purged with an inert gas. Timosaponin AI is soluble in the organic solvent DMSO at a concentration of approximately 10 mM.

Description

Timosaponin AI is a steroidal saponin that has been found in *A. asphodeloides* and has diverse biological activities.¹⁻³ It is also a metabolite of timosaponin AIII (Item No. 22236).⁴ Timosaponin AI inhibits 5-lipoxygenase (5-LO), COX-2, and dipeptidyl peptidase 4 (DPP-4; IC₅₀s = 3.29, 36.43, and 33.25 μ M, respectively).^{1,2} It reduces total cholesterol and triglyceride levels in 3T3-L1 differentiated adipocytes and increases glucose consumption in insulin-resistant BNL CL.2 mouse embryonic liver cells stimulated with insulin.² Timosaponin AI is cytotoxic to NRK rat kidney epithelial and MOLM-13 acute myeloid leukemia (AML) cells (EC₅₀s = 0.12 and 0.22 mM, respectively).³

References

1. Xie, L., Lee, D.Y.-W., Shang, Y., et al. Characterization of spirostanol glycosides and furostanol glycosides from anemarrhenae rhizoma as dual targeted inhibitors of 5-lipoxygenase and Cyclooxygenase-2 by employing a combination of affinity ultrafiltration and HPLC/MS. *Phytomedicine* **77**, 153284 (2020).
2. Jiang, S., Wu, X., Wang, Y., et al. The potential DPP-4 inhibitors from Xiao-Ke-An improve the glucolipid metabolism via the activation of AKT/GSK-3 β pathway. *Eur. J. Pharmacol.* **882**, 173272 (2020).
3. Carpineteyro Díaz, A.E., Herfindal, L., Rathe, B.A., et al. Cytotoxic saponins and other natural products from flowering tops of *Narthecium ossifragum* L. *Phytochemistry* **164**, 67-77 (2019).
4. Sun, Y., Liu, L., Peng, Y., et al. Metabolites characterization of timosaponin AIII *in vivo* and *in vitro* by using liquid chromatography-mass spectrometry. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **997**, 236-243 (2015).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM