

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Galactaric Acid

Item No. 36206

CAS Registry No.: 526-99-8

meso-Galactaric Acid, Mucic Acid, NSC 8127 Synonyms:

MF: $C_6H_{20}O_8$ FW: 210.1 **Purity:** ≥95% Supplied as: A solid -20°C Storage: Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Galactaric acid is supplied as a solid. A stock solution may be made by dissolving the galactaric acid in the solvent of choice, which should be purged with an inert gas. Galactaric acid is soluble in 1 M NaOH.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of galactaric acid can be prepared by directly dissolving the solid in aqueous buffers. The solubility of galactaric acid in PBS (pH 7.2) is approximately 1 mg/ml (sonicated). We do not recommend storing the aqueous solution for more than one day.

Description

Galactaric acid is a monosaccharide that has been found in grape musts and wine and an oxidized form of galacturonic acid.²⁻⁴ Galactaric acid (5-10 μM) increases mRNA expression of Runx2 in C3H/10T1/2 mouse mesenchymal stem cells.1 It increases alkaline phosphatase (ALP) levels in, and mineralization of, C3H/10T1/2 cells when used in combination with gelatin as a coating on a 3D-poly (lactic acid) (PLA) scaffold at concentrations ranging from 10 to 20 μM.

References

- 1. Ashwin, B., Abinaya, B., Prasith, T.P., et al. 3D-poly (lactic acid) scaffolds coated with gelatin and mucic acid for bone tissue engineering. Int. J. Biol. Macromol. 162, 523-532 (2020).
- Barth, D. and Wiebe, M.G. Enhancing fungal production of galactaric acid. Appl. Microbiol. Biotechnol. 101(10), 4033-4040 (2017).
- 3. Rautiainen, S., Lehtinen, P., Chen, J., et al. Selective oxidation of uronic acids into aldaric acids over gold catalyst. RSC Adv. 25, 19502-19507 (2015).
- 4. Žulj, M.M., Puhelek, I., Korenika, A.M.J., et al. Organic acid composition in Croatian predicate wines. Agriculturae Conspectus Scientificus 80, 113-117 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/19/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM